

$K^*(1410)$

$$I(J^P) = \frac{1}{2}(1^-)$$

 $K^*(1410)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
1414 ± 15 OUR AVERAGE		Error includes scale factor of 1.3.				
1380 ± 21 ± 19		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1420 ± 7 ± 10		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1437 ± 8 ± 16	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
1426 ± 8 ± 24	190k	² AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
1276 ⁺⁷² ₋₇₇		^{3,4} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
1367 ± 54		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1474 ± 25		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
1500 ± 30		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

¹ Using a parametrization for the $K\pi$ S -wave similar to ASTON 88 with fixed resonance width.

² Using a $K\pi$ S -wave parametrization with resonant and non-resonant contributions.

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

 $K^*(1410)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
232 ± 21 OUR AVERAGE		Error includes scale factor of 1.1.				
176 ± 52 ± 22		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
240 ± 18 ± 12		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
210 ± 20 ± 60	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
270 ± 20 ± 40	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
198 ⁺⁶¹ ₋₈₇		^{2,3} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
114 ± 101		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
275 ± 65		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
500 ± 100		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

¹ Using a $K\pi$ S -wave parametrization with resonant and non-resonant contributions.

² From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

³ Systematic uncertainties not estimated.

$K^*(1410)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K^*(892)\pi$	> 40 %	95%
Γ_2 $K\pi$	(6.6 ± 1.3) %	
Γ_3 $K\rho$	< 7 %	95%
Γ_4 γK^0	< 2.3 $\times 10^{-4}$	90%

 $K^*(1410)$ PARTIAL WIDTHS

$\Gamma(\gamma K^0)$					Γ_4
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<52.9	90	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$	

 $K^*(1410)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$						Γ_3/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.17	95	ASTON	84	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$						Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.16	95	ASTON	84	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
$0.066 \pm 0.010 \pm 0.008$	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

 $K^*(1410)$ REFERENCES

AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
BOITO	09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin	
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP