

# b-baryon ADMIXTURE ( $\Lambda_b$ , $\Xi_b$ , $\Omega_b$ )

## b-baryon ADMIXTURE MEAN LIFE

Each measurement of the  $b$ -baryon mean life is an average over an admixture of various  $b$  baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different  $b$ -baryon mean life. More  $b$ -baryon flavor specific channels are not included in the measurement.

VALUE ( $10^{-12}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.218^{+0.130}_{-0.115} \pm 0.042$		<sup>1</sup> ABAZOV	07S D0	Repl. by ABAZOV 12U
$1.22^{+0.22}_{-0.18} \pm 0.04$		<sup>1</sup> ABAZOV	05C D0	Repl. by ABAZOV 07S
$1.16 \pm 0.20 \pm 0.08$		<sup>2</sup> ABREU	99W DLPH	$e^+e^- \rightarrow Z$
$1.19 \pm 0.14 \pm 0.07$		<sup>3</sup> ABREU	99W DLPH	$e^+e^- \rightarrow Z$
$1.14 \pm 0.08 \pm 0.04$		<sup>4</sup> ABREU	99W DLPH	$e^+e^- \rightarrow Z$
$1.11^{+0.19}_{-0.18} \pm 0.05$		<sup>5</sup> ABREU	99W DLPH	$e^+e^- \rightarrow Z$
$1.29^{+0.24}_{-0.22} \pm 0.06$		<sup>5</sup> ACKERSTAFF	98G OPAL	$e^+e^- \rightarrow Z$
$1.20 \pm 0.08 \pm 0.06$		<sup>6</sup> BARATE	98D ALEP	$e^+e^- \rightarrow Z$
$1.21 \pm 0.11$		<sup>5</sup> BARATE	98D ALEP	$e^+e^- \rightarrow Z$
$1.32 \pm 0.15 \pm 0.07$		<sup>7</sup> ABE	96M CDF	$p\bar{p}$ at 1.8 TeV
$1.46^{+0.22}_{-0.21} \pm 0.07_{-0.09}$		ABREU	96D DLPH	Repl. by ABREU 99W
$1.10^{+0.19}_{-0.17} \pm 0.09$		<sup>5</sup> ABREU	96D DLPH	$e^+e^- \rightarrow Z$
$1.16 \pm 0.11 \pm 0.06$		<sup>5</sup> AKERS	96 OPAL	$e^+e^- \rightarrow Z$
$1.27^{+0.35}_{-0.29} \pm 0.09$		ABREU	95S DLPH	Repl. by ABREU 99W
$1.05^{+0.12}_{-0.11} \pm 0.09$	290	BUSKULIC	95L ALEP	Repl. by BARATE 98D
$1.04^{+0.48}_{-0.38} \pm 0.10$	11	<sup>8</sup> ABREU	93F DLPH	Excess $\Lambda\mu^-$ , decay lengths
$1.05^{+0.23}_{-0.20} \pm 0.08$	157	<sup>9</sup> AKERS	93 OPAL	Excess $\Lambda\ell^-$ , decay lengths
$1.12^{+0.32}_{-0.29} \pm 0.16$	101	<sup>10</sup> BUSKULIC	92I ALEP	Excess $\Lambda\ell^-$ , impact parameters

<sup>1</sup> Measured mean life using fully reconstructed  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  decays.

<sup>2</sup> Measured using  $\Lambda\ell^-$  decay length.

<sup>3</sup> Measured using  $p\ell^-$  decay length.

<sup>4</sup> This ABREU 99W result is the combined result of the  $\Lambda\ell^-$ ,  $p\ell^-$ , and excess  $\Lambda\mu^-$  impact parameter measurements.

<sup>5</sup> Measured using  $\Lambda_c\ell^-$  and  $\Lambda\ell^+\ell^-$ .

<sup>6</sup> Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>7</sup> Measured using  $\Lambda_c\ell^-$ .

<sup>8</sup> ABREU 93F superseded by ABREU 96D.

<sup>9</sup> AKERS 93 superseded by AKERS 96.<sup>10</sup> BUSKULIC 92I superseded by BUSKULIC 95L.

## ***b*-baryon ADMIXTURE DECAY MODES**

### **( $\Lambda_b, \Xi_b, \Omega_b$ )**

These branching fractions are actually an average over weakly decaying *b*-baryons weighted by their production rates at the LHC, LEP, and Tevatron, branching ratios, and detection efficiencies. They scale with the *b*-baryon production fraction  $B(b \rightarrow b\text{-baryon})$ .

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note "Production and Decay of *b*-Flavored Hadrons."

For inclusive branching fractions, e.g.,  $B \rightarrow D^\pm \text{ anything}$ , the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $p\mu^- \bar{\nu}$ anything	$(5.8_{-2.0}^{+2.3})\%$
$\Gamma_2$ $p\ell \bar{\nu}_\ell$ anything	$(5.6 \pm 1.2)\%$
$\Gamma_3$ $p$ anything	$(70 \pm 22)\%$
$\Gamma_4$ $\Lambda \ell^- \bar{\nu}_\ell$ anything	$(3.8 \pm 0.6)\%$
$\Gamma_5$ $\Lambda \ell^+ \nu_\ell$ anything	$(3.2 \pm 0.8)\%$
$\Gamma_6$ $\Lambda$ anything	$(39 \pm 7)\%$
$\Gamma_7$ $\Xi^- \ell^- \bar{\nu}_\ell$ anything	$(6.6 \pm 1.6) \times 10^{-3}$

## ***b*-baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Omega_b$ ) BRANCHING RATIOS**

$\Gamma(p\mu^- \bar{\nu} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
$5.8_{-1.9}^{+2.2} \pm 0.8$	125	<sup>11</sup> ABREU	95s DLPH	$e^+e^- \rightarrow Z$	

<sup>11</sup> ABREU 95S reports  $[\Gamma(b\text{-baryon} \rightarrow p\mu^- \bar{\nu} \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0049 \pm 0.0011_{-0.0011}^{+0.0015}$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\ell \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE (%)		DOCUMENT ID	TECN	COMMENT	
$5.6 \pm 0.9 \pm 0.7$		<sup>12</sup> BARATE	98V ALEP	$e^+e^- \rightarrow Z$	

<sup>12</sup> BARATE 98V reports  $[\Gamma(b\text{-baryon} \rightarrow p\ell \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = (4.72 \pm 0.66 \pm 0.44) \times 10^{-3}$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho\ell\bar{\nu}_\ell \text{ anything})/\Gamma(\rho \text{ anything})$  $\Gamma_2/\Gamma_3$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>8.0±1.2±1.4</b>	BARATE	98V ALEP	$e^+e^- \rightarrow Z$

 $\Gamma(\Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

The values and averages in this section serve only to show what values result if one assumes our  $B(b \rightarrow b\text{-baryon})$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determine  $B(b \rightarrow b\text{-baryon})$  as described in the note on "Production and Decay of  $b$ -Flavored Hadrons."

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±0.6 OUR AVERAGE</b>				
3.9±0.5±0.5		<sup>13</sup> BARATE	98D ALEP	$e^+e^- \rightarrow Z$
3.5±0.4±0.5		<sup>14</sup> AKERS	96 OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
3.6±0.9±0.5	262	<sup>15</sup> ABREU	95S DLPH	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
7.3±1.4±1.0	290	<sup>16</sup> BUSKULIC	95L ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	157	<sup>17</sup> AKERS	93 OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
8.3±2.5±1.1	101	<sup>18</sup> BUSKULIC	92I ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

<sup>13</sup> BARATE 98D reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00326 \pm 0.00016 \pm 0.00039$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>14</sup> AKERS 96 reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00291 \pm 0.00023 \pm 0.00025$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>15</sup> ABREU 95S reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0030 \pm 0.0006 \pm 0.0004$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>16</sup> BUSKULIC 95L reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0061 \pm 0.0006 \pm 0.0010$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>17</sup> AKERS 93 superseded by AKERS 96.

<sup>18</sup> BUSKULIC 92I reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0070 \pm 0.0010 \pm 0.0018$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

 $\Gamma(\Lambda\ell^+\nu_\ell \text{ anything})/\Gamma(\Lambda \text{ anything})$  $\Gamma_5/\Gamma_6$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.0±1.2±0.8</b>	ABBIENDI	99L OPAL	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0±1.2±0.7	ACKERSTAFF	97N OPAL	Repl. by ABBIENDI 99L
-------------	------------	----------	-----------------------

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

**39 ± 7 OUR AVERAGE**42 ± 6 ± 5      19 ABBIENDI      99L OPAL      e<sup>+</sup>e<sup>-</sup> → Z27<sup>+15</sup><sub>-9</sub> ± 3      20 ABREU      95C DLPH      e<sup>+</sup>e<sup>-</sup> → Z

• • • We do not use the following data for averages, fits, limits, etc. • • •

47 ± 7 ± 6      21 ACKERSTAFF 97N OPAL      Repl. by ABBIENDI 99L

19 ABBIENDI 99L reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.035 \pm 0.0032 \pm 0.0035$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

20 ABREU 95C reports  $0.28^{+0.17}_{-0.12}$  from a measurement of  $[\Gamma(b\text{-baryon} \rightarrow \Lambda \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$  assuming  $B(\bar{b} \rightarrow b\text{-baryon}) = 0.08 \pm 0.02$ , which we rescale to our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

21 ACKERSTAFF 97N reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0393 \pm 0.0046 \pm 0.0037$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Xi^- \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

**6.6 ± 1.6 OUR AVERAGE**6.4 ± 1.6 ± 0.8      22 BUSKULIC      96T ALEP      Excess  $\Xi^- \ell^-$  over  $\Xi^- \ell^+$ 7.0 ± 2.8 ± 0.9      23 ABREU      95V DLPH      Excess  $\Xi^- \ell^-$  over  $\Xi^- \ell^+$ 

22 BUSKULIC 96T reports  $[\Gamma(b\text{-baryon} \rightarrow \Xi^- \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00054 \pm 0.00011 \pm 0.00008$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

23 ABREU 95V reports  $[\Gamma(b\text{-baryon} \rightarrow \Xi^- \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00059 \pm 0.00021 \pm 0.0001$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 **$b$ -baryon ADMIXTURE ( $\Lambda_b$ ,  $\Xi_b$ ,  $\Omega_b$ ) REFERENCES**

ABAZOV	12U	PR D85 112003	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07S	PRL 99 142001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05C	PRL 94 102001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	99L	EPJ C9 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	99W	EPJ C10 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98D	EPJ C2 197	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ABE	96M	PRL 77 1439	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96D	ZPHY C71 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)

ABREU	95C	PL B347 447	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	93F	PL B311 379	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	93	PL B316 435	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)

---