



$\Lambda_c^+$  Status: \*\*\*\*

The parity of the  $\Lambda_c^+$  is defined to be positive (as are the parities of the proton, neutron, and  $\Lambda$ ). The quark content is  $u d c$ . Results of an analysis of  $p K^- \pi^+$  decays (JEZABEK 92) are consistent with  $J = 1/2$ . Nobody doubts that the spin is indeed 1/2.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

## $\Lambda_c^+$ MASS

Our value in 2004,  $2284.9 \pm 0.6$  MeV, was the average of the measurements now filed below as “not used.” The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes  $\Sigma_c - \Lambda_c^+$  and  $\Lambda_c^{*+} - \Lambda_c^+$  mass-difference measurements, but this doesn’t affect the  $\Lambda_c^+$  mass. The new (in 2006)  $\Lambda_c^+$  mass simply pushes all those other masses higher.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2286.46 ± 0.14 OUR FIT</b>				
<b>2286.46 ± 0.14</b>	4891	<sup>1</sup> AUBERT,B	05S BABR	$\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2284.7 $\pm 0.6$ $\pm 0.7$	1134	AVERY	91 CLEO	Six modes
2281.7 $\pm 2.7$ $\pm 2.6$	29	ALVAREZ	90B NA14	$p K^- \pi^+$
2285.8 $\pm 0.6$ $\pm 1.2$	101	BARLAG	89 NA32	$p K^- \pi^+$
2284.7 $\pm 2.3$ $\pm 0.5$	5	AGUILAR-...	88B LEBC	$p K^- \pi^+$
2283.1 $\pm 1.7$ $\pm 2.0$	628	ALBRECHT	88C ARG	$p K^- \pi^+, p \bar{K}^0, \Lambda 3\pi$
2286.2 $\pm 1.7$ $\pm 0.7$	97	ANJOS	88B E691	$p K^- \pi^+$
2281 $\pm 3$	2	JONES	87 HBC	$p K^- \pi^+$
2283 $\pm 3$	3	BOSETTI	82 HBC	$p K^- \pi^+$
2290 $\pm 3$	1	CALICCHIO	80 HYBR	$p K^- \pi^+$

<sup>1</sup>AUBERT,B 05S uses low-Q  $\Lambda K_S^0 K^+$  and  $\Sigma^0 K_S^0 K^+$  decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the  $\Lambda_c^+$  mass.

## $\Lambda_c^+$ MEAN LIFE


Measurements with an error  $\geq 100 \times 10^{-15}$  s or with fewer than 20 events have been omitted from the Listings.

VALUE ( $10^{-15}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>202.4 ± 3.1 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
203.5 $\pm 1.7 \pm 1.4$	304k	<sup>1</sup> AAIJ	19AG LHCb	$\Lambda_c^+ \rightarrow p K^- \pi^+$
204.6 $\pm 3.4 \pm 2.5$	8034	LINK	02c FOCS	$\Lambda_c^+ \rightarrow p K^- \pi^+$

$198.1 \pm 7.0 \pm 5.6$	1630	KUSHNIR...	01	SELX	$\Lambda_c^+ \rightarrow p K^- \pi^+$
$179.6 \pm 6.9 \pm 4.4$	4749	MAHMOOD	01	CLE2	$e^+ e^- \approx \gamma(4S)$
$215 \pm 16 \pm 8$	1340	FRABETTI	93D	E687	$\gamma \text{Be}, \Lambda_c^+ \rightarrow p K^- \pi^+$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
180 $\pm 30 \pm 30$	29	ALVAREZ	90	NA14	$\gamma, \Lambda_c^+ \rightarrow p K^- \pi^+$
200 $\pm 30 \pm 30$	90	FRABETTI	90	E687	$\gamma \text{Be}, \Lambda_c^+ \rightarrow p K^- \pi^+$
196 $\begin{array}{l} +23 \\ -20 \end{array}$	101	BARLAG	89	NA32	$p K^- \pi^+ + \text{c.c.}$
220 $\pm 30 \pm 20$	97	ANJOS	88B	E691	$p K^- \pi^+ + \text{c.c.}$

<sup>1</sup> AAIJ 19AG reports  $[\Lambda_c^+ \text{ MEAN LIFE}] / [D^\pm \text{ MEAN LIFE}] = 0.1956 \pm 0.0010 \pm 0.0013$   
which we multiply by our best value  $D^\pm \text{ MEAN LIFE} = (1.040 \pm 0.007) \times 10^{-12} \text{ s}$ .  
Our first error is their experiment's error and our second error is the systematic error  
from using our best value.

WEIGHTED AVERAGE  
 $202.4 \pm 3.1$  (Error scaled by 1.7)



## $\Lambda_c^+$ DECAY MODES

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction  $\Lambda_c^+ \rightarrow p \bar{K}^*(892)^0$  seen in  $\Lambda_c^+ \rightarrow p K^- \pi^+$  has been multiplied up to include  $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$  decays.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

**Hadronic modes with a  $p$  or  $n$ :  $S = -1$  final states**

$\Gamma_1$	$p K_S^0$	( $1.59 \pm 0.08$ ) %	S=1.1
$\Gamma_2$	$p K^- \pi^+$	( $6.28 \pm 0.32$ ) %	S=1.4
$\Gamma_3$	$p \bar{K}^*(892)^0$	[a] ( $1.96 \pm 0.27$ ) %	
$\Gamma_4$	$\Delta(1232)^{++} K^-$	( $1.08 \pm 0.25$ ) %	
$\Gamma_5$	$\Lambda(1520) \pi^+$	[a] ( $2.2 \pm 0.5$ ) %	
$\Gamma_6$	$p K^- \pi^+$ nonresonant	( $3.5 \pm 0.4$ ) %	
$\Gamma_7$	$p K_S^0 \pi^0$	( $1.97 \pm 0.13$ ) %	S=1.1
$\Gamma_8$	$n K_S^0 \pi^+$	( $1.82 \pm 0.25$ ) %	
$\Gamma_9$	$p \bar{K}^0 \eta$	( $1.6 \pm 0.4$ ) %	
$\Gamma_{10}$	$p K_S^0 \pi^+ \pi^-$	( $1.60 \pm 0.12$ ) %	S=1.1
$\Gamma_{11}$	$p K^- \pi^+ \pi^0$	( $4.46 \pm 0.30$ ) %	S=1.5
$\Gamma_{12}$	$p K^*(892)^- \pi^+$	[a] ( $1.4 \pm 0.5$ ) %	
$\Gamma_{13}$	$p(K^- \pi^+)_{\text{nonresonant}} \pi^0$	( $4.6 \pm 0.8$ ) %	
$\Gamma_{14}$	$\Delta(1232) \bar{K}^*(892)$	seen	
$\Gamma_{15}$	$p K^- 2\pi^+ \pi^-$	( $1.4 \pm 0.9$ ) $\times 10^{-3}$	
$\Gamma_{16}$	$p K^- \pi^+ 2\pi^0$	( $1.0 \pm 0.5$ ) %	

**Hadronic modes with a  $p$ :  $S = 0$  final states**

$\Gamma_{17}$	$p \pi^0$	< $2.7 \times 10^{-4}$	CL=90%
$\Gamma_{18}$	$p \eta$	( $1.24 \pm 0.30$ ) $\times 10^{-3}$	
$\Gamma_{19}$	$p \omega(782)^0$	( $9 \pm 4$ ) $\times 10^{-4}$	
$\Gamma_{20}$	$p \pi^+ \pi^-$	( $4.61 \pm 0.28$ ) $\times 10^{-3}$	
$\Gamma_{21}$	$p f_0(980)$	[a] ( $3.5 \pm 2.3$ ) $\times 10^{-3}$	
$\Gamma_{22}$	$p 2\pi^+ 2\pi^-$	( $2.3 \pm 1.4$ ) $\times 10^{-3}$	
$\Gamma_{23}$	$p K^+ K^-$	( $1.06 \pm 0.06$ ) $\times 10^{-3}$	
$\Gamma_{24}$	$p \phi$	[a] ( $1.06 \pm 0.14$ ) $\times 10^{-3}$	
$\Gamma_{25}$	$p K^+ K^- \text{non-}\phi$	( $5.3 \pm 1.2$ ) $\times 10^{-4}$	
$\Gamma_{26}$	$p \phi \pi^0$	( $10 \pm 4$ ) $\times 10^{-5}$	
$\Gamma_{27}$	$p K^+ K^- \pi^0 \text{nonresonant}$	< $6.3 \times 10^{-5}$	CL=90%

**Hadronic modes with a hyperon:  $S = -1$  final states**

$\Gamma_{28}$	$\Lambda \pi^+$	( $1.30 \pm 0.07$ ) %	S=1.1
$\Gamma_{29}$	$\Lambda \pi^+ \pi^0$	( $7.1 \pm 0.4$ ) %	S=1.1
$\Gamma_{30}$	$\Lambda \rho^+$	< $6$ %	CL=95%
$\Gamma_{31}$	$\Lambda \pi^- 2\pi^+$	( $3.64 \pm 0.29$ ) %	S=1.4
$\Gamma_{32}$	$\Sigma(1385)^+ \pi^+ \pi^-$ , $\Sigma^{*+} \rightarrow \Lambda \pi^+$	( $1.0 \pm 0.5$ ) %	
$\Gamma_{33}$	$\Sigma(1385)^- 2\pi^+$ , $\Sigma^{*-} \rightarrow \Lambda \pi^-$	( $7.6 \pm 1.4$ ) $\times 10^{-3}$	
$\Gamma_{34}$	$\Lambda \pi^+ \rho^0$	( $1.5 \pm 0.6$ ) %	
$\Gamma_{35}$	$\Sigma(1385)^+ \rho^0$ , $\Sigma^{*+} \rightarrow \Lambda \pi^+$	( $5 \pm 4$ ) $\times 10^{-3}$	
$\Gamma_{36}$	$\Lambda \pi^- 2\pi^+ \text{nonresonant}$	< $1.1$ %	CL=90%
$\Gamma_{37}$	$\Lambda \pi^- \pi^0 2\pi^+ \text{total}$	( $2.3 \pm 0.8$ ) %	
$\Gamma_{38}$	$\Lambda \pi^+ \eta$	[a] ( $1.84 \pm 0.26$ ) %	
$\Gamma_{39}$	$\Sigma(1385)^+ \eta$	[a] ( $9.1 \pm 2.0$ ) $\times 10^{-3}$	

$\Gamma_{40}$	$\Lambda\pi^+\omega$	[a] ( 1.5 ± 0.5 ) %	
$\Gamma_{41}$	$\Lambda\pi^-\pi^02\pi^+$ , no $\eta$ or $\omega$	< 8 × 10 <sup>-3</sup>	CL=90%
$\Gamma_{42}$	$\Lambda K^+\bar{K}^0$	( 5.7 ± 1.1 ) × 10 <sup>-3</sup>	S=1.9
$\Gamma_{43}$	$\Xi(1690)^0K^+$ , $\Xi^{*0} \rightarrow \Lambda\bar{K}^0$	( 1.6 ± 0.5 ) × 10 <sup>-3</sup>	
$\Gamma_{44}$	$\Sigma^0\pi^+$	( 1.29 ± 0.07 ) %	S=1.1
$\Gamma_{45}$	$\Sigma^+\pi^0$	( 1.25 ± 0.10 ) %	
$\Gamma_{46}$	$\Sigma^+\eta$	( 4.4 ± 2.0 ) × 10 <sup>-3</sup>	
$\Gamma_{47}$	$\Sigma^+\eta'$	( 1.5 ± 0.6 ) %	
$\Gamma_{48}$	$\Sigma^+\pi^+\pi^-$	( 4.50 ± 0.25 ) %	S=1.3
$\Gamma_{49}$	$\Sigma^+\rho^0$	< 1.7 %	CL=95%
$\Gamma_{50}$	$\Sigma^-2\pi^+$	( 1.87 ± 0.18 ) %	
$\Gamma_{51}$	$\Sigma^0\pi^+\pi^0$	( 3.5 ± 0.4 ) %	
$\Gamma_{52}$	$\Sigma^+\pi^0\pi^0$	( 1.55 ± 0.15 ) %	
$\Gamma_{53}$	$\Sigma^0\pi^-2\pi^+$	( 1.11 ± 0.30 ) %	
$\Gamma_{54}$	$\Sigma^+\pi^+\pi^-\pi^0$	—	
$\Gamma_{55}$	$\Sigma^+\omega$	[a] ( 1.70 ± 0.21 ) %	
$\Gamma_{56}$	$\Sigma^-\pi^02\pi^+$	( 2.1 ± 0.4 ) %	
$\Gamma_{57}$	$\Sigma^+K^+K^-$	( 3.5 ± 0.4 ) × 10 <sup>-3</sup>	S=1.1
$\Gamma_{58}$	$\Sigma^+\phi$	[a] ( 3.9 ± 0.6 ) × 10 <sup>-3</sup>	S=1.1
$\Gamma_{59}$	$\Xi(1690)^0K^+$ , $\Xi^{*0} \rightarrow \Sigma^+K^-$	( 1.02 ± 0.25 ) × 10 <sup>-3</sup>	
$\Gamma_{60}$	$\Sigma^+K^+K^-$ nonresonant	< 8 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{61}$	$\Xi^0K^+$	( 5.5 ± 0.7 ) × 10 <sup>-3</sup>	
$\Gamma_{62}$	$\Xi^-K^+\pi^+$	( 6.2 ± 0.6 ) × 10 <sup>-3</sup>	S=1.1
$\Gamma_{63}$	$\Xi(1530)^0K^+$	( 4.3 ± 0.9 ) × 10 <sup>-3</sup>	S=1.1

**Hadronic modes with a hyperon:  $S = 0$  final states**

$\Gamma_{64}$	$\Lambda K^+$	( 6.1 ± 1.2 ) × 10 <sup>-4</sup>	
$\Gamma_{65}$	$\Lambda K^+\pi^+\pi^-$	< 5 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{66}$	$\Sigma^0K^+$	( 5.2 ± 0.8 ) × 10 <sup>-4</sup>	
$\Gamma_{67}$	$\Sigma^0K^+\pi^+\pi^-$	< 2.6 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{68}$	$\Sigma^+K^+\pi^-$	( 2.1 ± 0.6 ) × 10 <sup>-3</sup>	
$\Gamma_{69}$	$\Sigma^+K^*(892)^0$	[a] ( 3.5 ± 1.0 ) × 10 <sup>-3</sup>	
$\Gamma_{70}$	$\Sigma^-K^+\pi^+$	< 1.2 × 10 <sup>-3</sup>	CL=90%

**Doubly Cabibbo-suppressed modes**

$\Gamma_{71}$	$pK^+\pi^-$	( 1.11 ± 0.18 ) × 10 <sup>-4</sup>
---------------	-------------	------------------------------------

**Semileptonic modes**

$\Gamma_{72}$	$\Lambda e^+\nu_e$	( 3.6 ± 0.4 ) %
$\Gamma_{73}$	$\Lambda\mu^+\nu_\mu$	( 3.5 ± 0.5 ) %

**Inclusive modes**

$\Gamma_{74}$	$e^+$ anything	( $3.95 \pm 0.35$ ) %
$\Gamma_{75}$	$p$ anything	( $50 \pm 16$ ) %
$\Gamma_{76}$	$n$ anything	( $50 \pm 16$ ) %
$\Gamma_{77}$	$\Lambda$ anything	( $38.2 \pm 2.4$ ) %
$\Gamma_{78}$	3prongs	( $24 \pm 8$ ) %

**$\Delta C = 1$  weak neutral current ( $C1$ ) modes, or  
Lepton Family number ( $LF$ ), or Lepton number ( $L$ ), or  
Baryon number ( $B$ ) violating modes**

$\Gamma_{79}$	$p e^+ e^-$	$C1$	< 5.5	$\times 10^{-6}$	CL=90%
$\Gamma_{80}$	$p \mu^+ \mu^-$ non-resonant	$C1$	< 7.7	$\times 10^{-8}$	CL=90%
$\Gamma_{81}$	$p e^+ \mu^-$	$LF$	< 9.9	$\times 10^{-6}$	CL=90%
$\Gamma_{82}$	$p e^- \mu^+$	$LF$	< 1.9	$\times 10^{-5}$	CL=90%
$\Gamma_{83}$	$\bar{p} 2e^+$	$L, B$	< 2.7	$\times 10^{-6}$	CL=90%
$\Gamma_{84}$	$\bar{p} 2\mu^+$	$L, B$	< 9.4	$\times 10^{-6}$	CL=90%
$\Gamma_{85}$	$\bar{p} e^+ \mu^+$	$L, B$	< 1.6	$\times 10^{-5}$	CL=90%
$\Gamma_{86}$	$\Sigma^- \mu^+ \mu^+$	$L$	< 7.0	$\times 10^{-4}$	CL=90%

[a] This branching fraction includes all the decay modes of the final-state resonance.

**CONSTRAINED FIT INFORMATION**

An overall fit to 41 branching ratios uses 62 measurements and one constraint to determine 21 parameters. The overall fit has a  $\chi^2 = 47.4$  for 42 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	54									
$x_7$	46	55								
$x_{10}$	44	64	39							
$x_{11}$	51	61	40	60						
$x_{28}$	54	66	44	42	43					
$x_{29}$	45	61	41	38	36	65				
$x_{31}$	51	37	28	41	60	45	36			
$x_{42}$	16	22	14	14	14	26	19	12		
$x_{44}$	51	55	38	37	40	74	58	44	20	
$x_{45}$	38	39	30	25	29	34	33	23	10	29
$x_{48}$	51	88	50	60	61	59	55	39	20	50
$x_{50}$	5	9	5	6	6	6	6	3	2	5
$x_{53}$	13	14	9	12	15	13	11	20	4	12
$x_{55}$	19	30	18	23	26	19	18	18	6	16
$x_{57}$	23	41	23	28	28	27	25	18	9	23
$x_{58}$	19	32	19	22	23	22	20	14	7	18
$x_{61}$	8	15	8	10	9	10	9	6	3	8
$x_{62}$	29	39	25	25	25	51	35	24	14	38
$x_{63}$	6	11	6	7	7	7	7	4	2	6
	$x_1$	$x_2$	$x_7$	$x_{10}$	$x_{11}$	$x_{28}$	$x_{29}$	$x_{31}$	$x_{42}$	$x_{44}$
$x_{48}$	36									
$x_{50}$	4	8								
$x_{53}$	7	14	1							
$x_{55}$	14	29	3	5						
$x_{57}$	17	45	4	6	13					
$x_{58}$	13	37	3	5	11	16				
$x_{61}$	6	13	1	2	5	6	5			
$x_{62}$	19	34	4	7	11	16	13	6		
$x_{63}$	4	10	1	2	3	4	4	2	4	
	$x_{45}$	$x_{48}$	$x_{50}$	$x_{53}$	$x_{55}$	$x_{57}$	$x_{58}$	$x_{61}$	$x_{62}$	

## $\Lambda_c^+$ BRANCHING RATIOS

A few really obsolete results have been omitted.

### — Hadronic modes with a $p$ : $S = -1$ final states —

#### $\Gamma(pK_S^0)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.59±0.08 OUR FIT</b>				Error includes scale factor of 1.1.
<b>1.52±0.08±0.03</b>	1243	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ , 4.599 GeV

#### $\Gamma(pK_S^0)/\Gamma(pK^-\pi^+)$

$\Gamma_1/\Gamma_2$

Measurements given as a  $\bar{K}^0$  ratio have been divided by 2 to convert to a  $K_S^0$  ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.254±0.012 OUR FIT</b>				Error includes scale factor of 1.4.
<b>0.234±0.020 OUR AVERAGE</b>				
0.23 ± 0.01 ± 0.02	1025	ALAM	98	CLE2 $e^+ e^- \approx \gamma(4S)$
0.22 ± 0.04 ± 0.03	133	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV
0.28 ± 0.09 ± 0.07	45	ANJOS	90	E691 $\gamma$ Be 70–260 GeV
0.31 ± 0.08 ± 0.02	73	ALBRECHT	88C	ARG $e^+ e^-$ 10 GeV

#### $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.28±0.32 OUR FIT</b>				Error includes scale factor of 1.4.
<b>6.3 ± 0.5 OUR AVERAGE</b>				Error includes scale factor of 2.0.
5.84±0.27±0.23	6.3k	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ , 4.599 GeV
6.84±0.24 <sup>+0.21</sup> <sub>-0.27</sub>	1.4k	<sup>1</sup> ZUPANC	14	BELL $e^+ e^- \rightarrow D^{(*)-} \bar{p}\pi^+$ recoil
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.0 ± 1.3		<sup>2</sup> PDG	02	See footnote

<sup>1</sup>This ZUPANC 14 value is the FIRST-EVER model-independent measurement of a  $\Lambda_c^+$  branching fraction.

<sup>2</sup>See the note by P. Burchat, " $\Lambda_c^+$  Branching Fractions," in any edition of the Review from 2002 through 2014 for how this value was obtained. It is now obsolete.

#### $\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$

$\Gamma_3/\Gamma_2$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.31±0.04 OUR AVERAGE</b>				
0.29±0.04±0.03		<sup>1</sup> AITALA	00	E791 $\pi^- N$ , 500 GeV
0.35 <sup>+0.06</sup> <sub>-0.07</sub> ±0.03	39	BOZEK	93	NA32 $\pi^- Cu$ 230 GeV
0.42±0.24	12	BASILE	81B	CNTR $pp \rightarrow \Lambda_c^+ e^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.35±0.11		BARLAG	90D	NA32 See BOZEK 93

<sup>1</sup>AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38$   $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays.

$\Gamma(\Delta(1232)^{++} K^-)/\Gamma(pK^-\pi^+)$  $\Gamma_4/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.17±0.04 OUR AVERAGE</b>	Error includes scale factor of 1.1.			
0.18±0.03±0.03	1	AITALA 00	E791	$\pi^- N$ , 500 GeV
0.12 <sup>+0.04</sup> <sub>-0.05</sub> ±0.05	14	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV
0.40±0.17	17	BASILE 81B	CNTR	$p p \rightarrow \Lambda_c^+ e^- X$

<sup>1</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38$   $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays.

 $\Gamma(\Lambda(1520)\pi^+)/\Gamma(pK^-\pi^+)$  $\Gamma_5/\Gamma_2$ Unseen decay modes of the  $\Lambda(1520)$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.35±0.08 OUR AVERAGE</b>				
0.34±0.08±0.05	1	AITALA 00	E791	$\pi^- N$ , 500 GeV
0.40 <sup>+0.18</sup> <sub>-0.13</sub> ±0.09	12	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV

<sup>1</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38$   $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays.

 $\Gamma(pK^-\pi^+ \text{ nonresonant})/\Gamma(pK^-\pi^+)$  $\Gamma_6/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.55±0.06 OUR AVERAGE</b>				
0.55±0.06±0.04	1	AITALA 00	E791	$\pi^- N$ , 500 GeV
0.56 <sup>+0.07</sup> <sub>-0.09</sub> ±0.05	71	BOZEK 93	NA32	$\pi^- Cu$ 230 GeV

<sup>1</sup> AITALA 00 makes a coherent 5-dimensional amplitude analysis of  $946 \pm 38$   $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays.

 $\Gamma(pK_S^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.97±0.13 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>1.87±0.13±0.05</b>	558	ABLIKIM 16	BES3	$e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(pK_S^0\pi^0)/\Gamma(pK^-\pi^+)$  $\Gamma_7/\Gamma_2$ Measurements given as a  $\bar{K}^0$  ratio have been divided by 2 to convert to a  $K_S^0$  ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.314±0.018 OUR FIT</b>				
<b>0.33 ±0.03 ±0.04</b>	774	ALAM 98	CLE2	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(nK_S^0\pi^+)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.82±0.23±0.11</b>	83	ABLIKIM 17H	BES3	$e^+ e^-$ at 4.6 GeV

 $\Gamma(p\bar{K}^0\eta)/\Gamma(pK^-\pi^+)$  $\Gamma_9/\Gamma_2$ Unseen decay modes of the  $\eta$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.25±0.04±0.04</b>	57	AMMAR 95	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(pK_S^0\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.60±0.12 OUR FIT</b>		Error includes scale factor of 1.1.		
<b>1.53±0.11±0.09</b>	485	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(pK_S^0\pi^+\pi^-)/\Gamma(pK^-\pi^+)$   $\Gamma_{10}/\Gamma_2$ Measurements given as a  $\bar{K}^0$  ratio have been divided by 2 to convert to a  $K_S^0$  ratio.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.255±0.015 OUR FIT</b>		Error includes scale factor of 1.1.		
<b>0.257±0.031 OUR AVERAGE</b>				
0.26 ± 0.02 ± 0.03	985	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$
0.22 ± 0.06 ± 0.02	83	AVERY	91	CLEO $e^+e^-$ 10.5 GeV
0.49 ± 0.18 ± 0.04	12	BARLAG	90D	NA32 $\pi^-$ 230 GeV

 $\Gamma(pK^-\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.46±0.30 OUR FIT</b>		Error includes scale factor of 1.5.		
<b>4.53±0.23±0.30</b>	1849	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(pK^-\pi^+\pi^0)/\Gamma(pK^-\pi^+)$   $\Gamma_{11}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.71 ± 0.04 OUR FIT</b>		Error includes scale factor of 2.4.		
<b>0.685±0.019 OUR AVERAGE</b>				
0.685±0.007±0.018	242k	PAL	17	BELL $e^+e^- \approx \gamma(4S), \gamma(5S)$
0.67 ± 0.04 ± 0.11	2.6k	ALAM	98	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(pK^*(892)^-\pi^+)/\Gamma(pK_S^0\pi^+\pi^-)$   $\Gamma_{12}/\Gamma_{10}$ Unseen decay modes of the  $K^*(892)^-$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.88±0.28</b>	17	ALEEV	94	BIS2 $nN$ 20–70 GeV

 $\Gamma(p(K^-\pi^+)_{\text{nonresonant}}\pi^0)/\Gamma(pK^-\pi^+)$   $\Gamma_{13}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.73±0.12±0.05</b>	67	BOZEK	93	NA32 $\pi^-$ Cu 230 GeV

 $\Gamma(\Delta(1232)\bar{K}^*(892))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	35	AMENDOLIA	87	SPEC $\gamma$ Ge-Si

 $\Gamma(pK^-\pi^+\pi^-)/\Gamma(pK^-\pi^+)$   $\Gamma_{15}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.022±0.015</b>		BARLAG	90D	NA32 $\pi^-$ 230 GeV

 $\Gamma(pK^-\pi^+2\pi^0)/\Gamma(pK^-\pi^+)$   $\Gamma_{16}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.16±0.07±0.03</b>	15	BOZEK	93	NA32 $\pi^-$ Cu 230 GeV

**Hadronic modes with a  $p$ :  $S = 0$  final states** **$\Gamma(p\pi^0)/\Gamma_{\text{total}}$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{17}/\Gamma$
$<2.7 \times 10^{-4}$	90	ABLIKIM	17Q	BES3 $e^+ e^-$ at 4.6 GeV	

 **$\Gamma(p\eta)/\Gamma_{\text{total}}$** Unseen decay modes of the  $\eta$  are included.

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{18}/\Gamma$
$1.24 \pm 0.28 \pm 0.10$	52	ABLIKIM	17Q	BES3 $\eta \rightarrow 2\gamma, \pi^+\pi^0\pi^-$	

 **$\Gamma(p\omega(782)^0)/\Gamma_{\text{total}}$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{19}/\Gamma$
$9.4 \pm 3.2 \pm 2.2$	13	AAIJ	18N	LHCb    Seen in $\Lambda_c^+ \rightarrow p\mu^+\mu^-$	

 **$\Gamma(p\pi^+\pi^-)/\Gamma(pK^-\pi^+)$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{20}/\Gamma_2$
<b><math>7.35 \pm 0.24</math> OUR AVERAGE</b>				Error includes scale factor of 1.3.	
7.44 $\pm 0.08 \pm 0.18$	20k	AAIJ	18V	LHCb $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$	
6.70 $\pm 0.48 \pm 0.25$	495	ABLIKIM	16U	BES3 $e^+ e^-$ at 4.599 GeV	
6.9 $\pm 3.6$	5	BARLAG	90D	NA32 $\pi^-$ 230 GeV	

 **$\Gamma(pf_0(980))/\Gamma(pK^-\pi^+)$** Unseen decay modes of the  $f_0(980)$  are included.

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{21}/\Gamma_2$
<b>0.055 <math>\pm 0.036</math></b>	BARLAG	90D	NA32 $\pi^-$ 230 GeV	

 **$\Gamma(p2\pi^+2\pi^-)/\Gamma(pK^-\pi^+)$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{22}/\Gamma_2$
<b>0.036 <math>\pm 0.023</math></b>	BARLAG	90D	NA32 $\pi^-$ 230 GeV	

 **$\Gamma(pK^+K^-)/\Gamma(pK^-\pi^+)$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23}/\Gamma_2$
<b><math>1.70 \pm 0.04</math> OUR AVERAGE</b>					
1.70 $\pm 0.03 \pm 0.03$	3.4k	AAIJ	18V	LHCb $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$	
1.4 $\pm 0.2 \pm 0.2$	676	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$	
3.9 $\pm 0.9 \pm 0.7$	214	ALEXANDER	96C	CLE2 $e^+ e^- \approx \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
9.6 $\pm 2.9 \pm 1.0$	30	FRABETTI	93H	E687 $\gamma Be, \bar{E}_\gamma$ 220 GeV	
4.8 $\pm 2.7$		BARLAG	90D	NA32 $\pi^-$ 230 GeV	

 **$\Gamma(p\phi)/\Gamma(pK^-\pi^+)$** Unseen decay modes of the  $\phi$  are included.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{24}/\Gamma_2$
<b><math>1.70 \pm 0.21</math> OUR AVERAGE</b>					
1.81 $\pm 0.33 \pm 0.13$	44	ABLIKIM	16U	BES3 $e^+ e^-$ at 4.599 GeV	
1.5 $\pm 0.2 \pm 0.2$	345	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$	
2.4 $\pm 0.6 \pm 0.3$	54	ALEXANDER	96C	CLE2 $e^+ e^- \approx \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.0 $\pm 2.7$		BARLAG	90D	NA32 $\pi^-$ 230 GeV	

$\Gamma(pK^+K^-\text{non-}\phi)/\Gamma(pK^-\pi^+)$  $\Gamma_{25}/\Gamma_2$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.4 ± 1.8 OUR AVERAGE</b>				
9.36 ± 2.22 ± 0.71	38	ABLIKIM	16U	BES3 $e^+e^-$ at 4.599 GeV
7 ± 2 ± 2	344	ABE	02C	BELL $e^+e^- \approx \gamma(4S)$

 $\Gamma(p\phi\pi^0)/\Gamma(pK^-\pi^+)$  $\Gamma_{26}/\Gamma_2$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.538 ± 0.641 ± 0.077</b>	PAL	17	BELL $e^+e^- \approx \gamma(4S), \gamma(5S)$

 $\Gamma(pK^+K^-\pi^0\text{nonresonant})/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;6.3 × 10<sup>-5</sup></b>	90	PAL	17	BELL $e^+e^- \approx \gamma(4S), \gamma(5S)$

**Hadronic modes with a hyperon:  $S = -1$  final states** $\Gamma(\Lambda\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{28}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.30 ± 0.07 OUR FIT</b>				Error includes scale factor of 1.1.
<b>1.24 ± 0.07 ± 0.03</b>	706	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$  $\Gamma_{28}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.207 ± 0.009 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.204 ± 0.019 OUR AVERAGE</b>				
0.217 ± 0.013 ± 0.020	750	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.18 ± 0.03 ± 0.04		ALBRECHT	92	ARG $e^+e^- \approx 10.4$ GeV
0.18 ± 0.03 ± 0.03	87	AVERY	91	CLEO $e^+e^-$ 10.5 GeV

 $\Gamma(\Lambda\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{29}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.1 ± 0.4 OUR FIT</b>				Error includes scale factor of 1.1.
<b>7.01 ± 0.37 ± 0.19</b>	1497	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(\Lambda\pi^+\pi^0)/\Gamma(pK^-\pi^+)$  $\Gamma_{29}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.12 ± 0.05 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.73 ± 0.09 ± 0.16</b>	464	AVERY	94	CLE2 $e^+e^- \approx \gamma(3S), \gamma(4S)$

 $\Gamma(\Lambda\rho^+)/\Gamma(pK^-\pi^+)$  $\Gamma_{30}/\Gamma_2$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.95</b>	95	AVERY	94	CLE2 $e^+e^- \approx \gamma(3S), \gamma(4S)$

 $\Gamma(\Lambda\pi^-2\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.64 ± 0.29 OUR FIT</b>				Error includes scale factor of 1.4.
<b>3.81 ± 0.24 ± 0.18</b>	609	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

$\Gamma(\Lambda\pi^- 2\pi^+)/\Gamma(pK^-\pi^+)$  $\Gamma_{31}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.58 ± 0.04 OUR FIT</b>	Error includes scale factor of 1.9.			
<b>0.522 ± 0.032 OUR AVERAGE</b>				
0.508 ± 0.024 ± 0.024	1356	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.65 ± 0.11 ± 0.12	289	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV
0.82 ± 0.29 ± 0.27	44	ANJOS	90	E691 $\gamma$ Be 70–260 GeV
0.94 ± 0.41 ± 0.13	10	BARLAG	90D	NA32 $\pi^-$ 230 GeV
0.61 ± 0.16 ± 0.04	105	ALBRECHT	88C	ARG $e^+ e^-$ 10 GeV

 $\Gamma(\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^- 2\pi^+)$  $\Gamma_{32}/\Gamma_{31}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.28 ± 0.10 ± 0.08</b>	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Sigma(1385)^-2\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-)/\Gamma(\Lambda\pi^- 2\pi^+)$  $\Gamma_{33}/\Gamma_{31}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.21 ± 0.03 ± 0.02</b>	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^+\rho^0)/\Gamma(\Lambda\pi^- 2\pi^+)$  $\Gamma_{34}/\Gamma_{31}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.40 ± 0.12 ± 0.12</b>	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^- 2\pi^+)$  $\Gamma_{35}/\Gamma_{31}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.14 ± 0.09 ± 0.07</b>	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^- 2\pi^+ \text{nonresonant})/\Gamma(\Lambda\pi^- 2\pi^+)$  $\Gamma_{36}/\Gamma_{31}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.3</b>	90	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

 $\Gamma(\Lambda\pi^- \pi^0 2\pi^+ \text{total})/\Gamma(pK^-\pi^+)$  $\Gamma_{37}/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.36 ± 0.09 ± 0.09</b>	50	<sup>1</sup> CRONIN-HEN..03	CLE3	$e^+ e^- \approx \gamma(4S)$

<sup>1</sup> CRONIN-HENNESSY 03 finds this channel to be dominantly  $\Lambda\eta\pi^+$  and  $\Lambda\omega\pi^+$ ; see below.

 $\Gamma(\Lambda\pi^+\eta)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.84 ± 0.21 ± 0.15</b>	154	ABLIKIM	19Y	BES3 $e^+ e^-$ at 4.6 GeV

 $\Gamma(\Lambda\pi^+\eta)/\Gamma(pK^-\pi^+)$  $\Gamma_{38}/\Gamma_2$ Unseen decay modes of the  $\eta$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.36 ± 0.07 OUR AVERAGE</b>				
0.41 ± 0.17 ± 0.10	11	CRONIN-HEN..03	CLE3	$e^+ e^- \approx \gamma(4S)$
0.35 ± 0.05 ± 0.06	116	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$

$\Gamma(\Sigma(1385)^+\eta)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

<u>VALUE</u> (units $10^{-2}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.91±0.18±0.09</b>	54	ABLIKIM	19Y	BES3 $e^+ e^-$ at 4.6 GeV	

 $\Gamma(\Sigma(1385)^+\eta)/\Gamma(pK^-\pi^+)$   $\Gamma_{39}/\Gamma_2$ Unseen decay modes of the  $\Sigma(1385)^+$  and  $\eta$  are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.17±0.04±0.03</b>	54	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Lambda\pi^+\omega)/\Gamma(pK^-\pi^+)$   $\Gamma_{40}/\Gamma_2$ Unseen decay modes of the  $\omega$  are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.24±0.06±0.06</b>	32	CRONIN-HEN..03	CLE3	$e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Lambda\pi^-\pi^02\pi^+, \text{no } \eta \text{ or } \omega)/\Gamma(pK^-\pi^+)$   $\Gamma_{41}/\Gamma_2$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>&lt;0.13</b>	90	CRONIN-HEN..03	CLE3	$e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(pK^-\pi^+)$   $\Gamma_{42}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.090±0.017 OUR FIT</b>				Error includes scale factor of 1.9.	
<b>0.131±0.020 OUR AVERAGE</b>					
0.142±0.018±0.022	251	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
0.12 ± 0.02 ± 0.02	59	AMMAR	95	CLE2 $e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Xi(1690)^0 K^+, \Xi^* \rightarrow \Lambda \bar{K}^0)/\Gamma(\Lambda K^+\bar{K}^0)$   $\Gamma_{43}/\Gamma_{42}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.28±0.07 OUR AVERAGE</b>					
0.32±0.10±0.04	84±24	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
0.26±0.08±0.03	93	ABE	02C	BELL $e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(\Lambda\pi^+)$   $\Gamma_{42}/\Gamma_{28}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.44 ± 0.08 OUR FIT</b>				Error includes scale factor of 2.0.	
<b>0.395±0.026±0.036</b>	460 ± 30	AUBERT	07U	BABR $e^+ e^- \approx \gamma(4S)$	

 $\Gamma(\Sigma^0\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>1.29±0.07 OUR FIT</b>				Error includes scale factor of 1.1.	
<b>1.27±0.08±0.03</b>	522	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ , 4.599 GeV	

 $\Gamma(\Sigma^0\pi^+)/\Gamma(pK^-\pi^+)$   $\Gamma_{44}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<b> </b>
<b>0.206±0.010 OUR FIT</b>				Error includes scale factor of 1.2.	
<b>0.20 ± 0.04 OUR AVERAGE</b>					
0.21 ± 0.02 ± 0.04	196	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$	
0.17 ± 0.06 ± 0.04		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV	

$\Gamma(\Sigma^0\pi^+)/\Gamma(\Lambda\pi^+)$  $\Gamma_{44}/\Gamma_{28}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.99 ± 0.04 OUR FIT</b>				
<b>0.98 ± 0.05 OUR AVERAGE</b>				

0.977 ± 0.015 ± 0.051      33k      AUBERT      07U      BABR       $e^+e^- \approx \gamma(4S)$   
 1.09 ± 0.11 ± 0.19      750      LINK      05F      FOCS       $\gamma$  nucleus,  $\bar{E}_\gamma \approx 180$  GeV

 $\Gamma(\Sigma^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.25 ± 0.10 OUR FIT</b>				
<b>1.18 ± 0.10 ± 0.03</b>	309	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(\Sigma^+\pi^0)/\Gamma(pK^-\pi^+)$  $\Gamma_{45}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.199 ± 0.015 OUR FIT</b>				
<b>0.20 ± 0.03 ± 0.03</b>	93	KUBOTA	93	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+\eta)/\Gamma(pK^-\pi^+)$  $\Gamma_{46}/\Gamma_2$ Unseen decay modes of the  $\eta$  are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.11 ± 0.03 ± 0.02</b>	26	AMMAR	95	CLE2 $e^+e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+\eta)/\Gamma(\Sigma^+\pi^0)$  $\Gamma_{46}/\Gamma_{45}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.35 ± 0.16 ± 0.02</b>	15	<sup>1</sup> ABLIKIM	19X	BES3 $e^+e^-$ at 4.6 GeV

<sup>1</sup> ABLIKIM 19X report evidence for the observation of the decay  $\Lambda_c^+ \rightarrow \Sigma^+ \eta$  at  $2.5\sigma$  significance.

 $\Gamma(\Sigma^+\eta')/\Gamma(\Sigma^+\omega)$  $\Gamma_{47}/\Gamma_{55}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.86 ± 0.34 ± 0.04</b>	13	<sup>1</sup> ABLIKIM	19X	BES3 $e^+e^-$ at 4.6 GeV

<sup>1</sup> ABLIKIM 19X report evidence for the observation of the decay  $\Lambda_c^+ \rightarrow \Sigma^+ \eta'$  at  $3.2\sigma$  significance.

 $\Gamma(\Sigma^+\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{48}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.50 ± 0.25 OUR FIT</b>				Error includes scale factor of 1.3.
<b>4.25 ± 0.24 ± 0.20</b>	1156	ABLIKIM	16	BES3 $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c$ , 4.599 GeV

 $\Gamma(\Sigma^+\pi^+\pi^-)/\Gamma(pK^-\pi^+)$  $\Gamma_{48}/\Gamma_2$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.716 ± 0.019 OUR FIT</b>				
<b>0.720 ± 0.024 OUR AVERAGE</b>				

0.719 ± 0.003 ± 0.024	2.7M	BERGER	18	BELL $e^+e^- \approx \gamma(4S)$
0.74 ± 0.07 ± 0.09	487	KUBOTA	93	CLE2 $e^+e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72 ± 0.14	47 ± 9	VAZQUEZ-JA...08	SELX	$\Sigma^-$ nucleus, 600 GeV
0.54 ± 0.18	11	BARLAG	92	NA32 $\pi^-$ Cu 230 GeV

$\Gamma(\Sigma^+ \rho^0)/\Gamma(p K^- \pi^+)$					$\Gamma_{49}/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.27</b>	95	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$	
$\Gamma(\Sigma^- 2\pi^+)/\Gamma_{\text{total}}$					$\Gamma_{50}/\Gamma$
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.87 ± 0.18 OUR FIT</b>					
<b>1.81 ± 0.17 ± 0.09</b>	161	ABLIKIM	17Y	BES3 $e^+ e^-$ at 4.6 GeV	
$\Gamma(\Sigma^- 2\pi^+)/\Gamma(p K^- \pi^+)$					$\Gamma_{50}/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.297 ± 0.030 OUR FIT</b>		Error includes scale factor of 1.1.			
<b>0.314 ± 0.067</b>	30 ± 6	VAZQUEZ-JA...08	SELX	$\Sigma^-$ nucleus, 600 GeV	
$\Gamma(\Sigma^- 2\pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$					$\Gamma_{50}/\Gamma_{48}$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.42 ± 0.04 OUR FIT</b>		Error includes scale factor of 1.1.			
<b>0.53 ± 0.15 ± 0.07</b>	56	FRABETTI	94E	E687 $\gamma$ Be, $\bar{E}_\gamma$ 220 GeV	
$\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(p K^- \pi^+)$					$\Gamma_{51}/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.56 ± 0.05 OUR AVERAGE</b>		Error includes scale factor of 1.5.			
0.575 ± 0.005 ± 0.036	2.7M	BERGER	18	BELL $e^+ e^- \approx \gamma(4S)$	
0.36 ± 0.09 ± 0.10	117	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$	
$\Gamma(\Sigma^+ \pi^0 \pi^0)/\Gamma(p K^- \pi^+)$					$\Gamma_{52}/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.247 ± 0.006 ± 0.019</b>	925k	BERGER	18	BELL $e^+ e^- \approx \gamma(4S)$	
$\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(p K^- \pi^+)$					$\Gamma_{53}/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.18 ± 0.05 OUR FIT</b>					
<b>0.21 ± 0.05 ± 0.05</b>	90	AVERY	94	CLE2 $e^+ e^- \approx \gamma(3S), \gamma(4S)$	
$\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(\Lambda \pi^- 2\pi^+)$					$\Gamma_{53}/\Gamma_{31}$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.31 ± 0.08 OUR FIT</b>					
<b>0.26 ± 0.06 ± 0.09</b>	480	LINK	05F	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
$\Gamma(\Sigma^+ \omega)/\Gamma_{\text{total}}$					$\Gamma_{55}/\Gamma$
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.70 ± 0.21 OUR FIT</b>					
<b>1.56 ± 0.20 ± 0.07</b>	157	ABLIKIM	16	BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ , 4.599 GeV	
$\Gamma(\Sigma^+ \omega)/\Gamma(p K^- \pi^+)$					$\Gamma_{55}/\Gamma_2$
Unseen decay modes of the $\omega$ are included.					
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.271 ± 0.031 OUR FIT</b>					
<b>0.54 ± 0.13 ± 0.06</b>	107	KUBOTA	93	CLE2 $e^+ e^- \approx \gamma(4S)$	

$\Gamma(\Sigma^-\pi^0 2\pi^+)/\Gamma_{\text{total}}$ 

VALUE (%)	EVTS
<b>2.11±0.33±0.14</b>	88

 $\Gamma_{56}/\Gamma$ 

DOCUMENT ID	TECN	COMMENT
ABLIKIM	17Y BES3	$e^+ e^-$ at 4.6 GeV

 $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(p K^- \pi^+)$ 

VALUE	EVTS
<b>0.056±0.006 OUR FIT</b>	
<b>0.070±0.011±0.011</b>	59

 $\Gamma_{57}/\Gamma_2$ 

DOCUMENT ID	TECN	COMMENT
AVERY	93 CLE2	$e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ 

VALUE	EVTS
<b>0.078±0.008 OUR FIT</b>	

**0.074±0.009 OUR AVERAGE**

VALUE	EVTS
0.076±0.007±0.009	246
0.071±0.011±0.011	103

 $\Gamma_{57}/\Gamma_{48}$ 

DOCUMENT ID	TECN	COMMENT
ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$
LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

 $\Gamma(\Sigma^+ \phi)/\Gamma(p K^- \pi^+)$ Unseen decay modes of the  $\phi$  are included.

VALUE	EVTS
<b>0.062±0.009 OUR FIT</b>	

VALUE	EVTS
<b>0.069±0.023±0.016</b>	26

 $\Gamma_{58}/\Gamma_2$ 

DOCUMENT ID	TECN	COMMENT
AVERY	93 CLE2	$e^+ e^- \approx 10.5$ GeV

 $\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Unseen decay modes of the  $\phi$  are included.

VALUE	EVTS
<b>0.087±0.012 OUR FIT</b>	

VALUE	EVTS
<b>0.086±0.012 OUR AVERAGE</b>	

VALUE	EVTS
0.085±0.012±0.012	129
0.087±0.016±0.006	57

 $\Gamma_{58}/\Gamma_{48}$ 

DOCUMENT ID	TECN	COMMENT
ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$
LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

 $\Gamma(\Xi(1690)^0 K^+, \Xi^*0 \rightarrow \Sigma^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$  $\Gamma_{59}/\Gamma_{48}$ 

VALUE	EVTS
<b>0.023±0.005 OUR AVERAGE</b>	

VALUE	EVTS
0.023±0.005±0.005	75

VALUE	EVTS
0.022±0.006±0.006	34

 $\Gamma_{59}/\Gamma_{48}$ 

DOCUMENT ID	TECN	COMMENT
ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$
LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

 $\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant})/\Gamma(\Sigma^+ \pi^+ \pi^-)$  $\Gamma_{60}/\Gamma_{48}$ 

VALUE	CL%
<b>&lt;0.018</b>	90

VALUE	CL%
<0.028	90

DOCUMENT ID	TECN	COMMENT
ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$
LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

 $\Gamma(\Xi^0 K^+)/\Gamma_{\text{total}}$  $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS
<b>5.5 ± 0.7 OUR FIT</b>	

VALUE	EVTS
<b>5.90±0.86±0.39</b>	68

DOCUMENT ID	TECN	COMMENT
ABLIKIM	18Y BES3	$e^+ e^-$ at 4.6 GeV

 $\Gamma(\Xi^0 K^+)/\Gamma(p K^- \pi^+)$  $\Gamma_{61}/\Gamma_2$ 

VALUE	EVTS
<b>0.088±0.012 OUR FIT</b>	

VALUE	EVTS
<b>0.078±0.013±0.013</b>	56


DOCUMENT ID	TECN	COMMENT
AVERY	93 CLE2	$e^+ e^- \approx 10.5$ GeV

### $\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$

$\Gamma_{62}/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.099±0.009 OUR FIT</b>				
<b>0.098±0.021 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
0.14 ± 0.03 ± 0.02	34	ALBRECHT	95B	ARG $e^+ e^- \approx 10.4$ GeV
0.079±0.013±0.014	60	AVERY	93	CLE2 $e^+ e^- \approx 10.5$ GeV
0.15 ± 0.04 ± 0.03	30	AVERY	91	CLEO $e^+ e^- 10.5$ GeV

WEIGHTED AVERAGE  
0.098±0.021 (Error scaled by 1.3)



$$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(p K^- \pi^+)$$

### $\Gamma(\Xi^- K^+ \pi^+)/\Gamma(\Lambda \pi^+)$

$\Gamma_{62}/\Gamma_{28}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.48 ± 0.04 OUR FIT</b>				
<b>0.480±0.016±0.039</b>	2665 ± 84	AUBERT	07U BABR	$e^+ e^- \approx \Upsilon(4S)$

### $\Gamma(\Xi(1530)^0 K^+)/\Gamma_{\text{total}}$

$\Gamma_{63}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.3 ± 0.9 OUR FIT</b>				Error includes scale factor of 1.1.
<b>5.02±0.99±0.31</b>	60	ABLIKIM	18Y BES3	$e^+ e^-$ at 4.6 GeV

### $\Gamma(\Xi(1530)^0 K^+)/\Gamma(p K^- \pi^+)$

$\Gamma_{63}/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.068±0.014 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.053±0.016±0.010</b>	24	AVERY	93 CLE2	$e^+ e^- \approx 10.5$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05 ± 0.02 ± 0.01	11	ALBRECHT	95B ARG	$e^+ e^- \approx 10.4$ GeV

**Hadronic modes with a hyperon:  $S = 0$  final states** $\Gamma(\Lambda K^+)/\Gamma(\Lambda \pi^+)$  $\Gamma_{64}/\Gamma_{28}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.047 \pm 0.009</math> OUR AVERAGE</b>	Error includes scale factor of 1.8.			
$0.044 \pm 0.004 \pm 0.003$	$1162 \pm 101$	AUBERT	07U BABR	$e^+ e^- \approx \gamma(4S)$
$0.074 \pm 0.010 \pm 0.012$	265	ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Lambda K^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+)$  $\Gamma_{65}/\Gamma_{28}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.1 \times 10^{-2}$	90	AUBERT	07U BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0 \pi^+)$  $\Gamma_{66}/\Gamma_{44}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.040 \pm 0.006</math> OUR AVERAGE</b>				
$0.038 \pm 0.005 \pm 0.003$	$366 \pm 52$	AUBERT	07U BABR	$e^+ e^- \approx \gamma(4S)$
$0.056 \pm 0.014 \pm 0.008$	75	ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^0 K^+ \pi^+ \pi^-)/\Gamma(\Sigma^0 \pi^+)$  $\Gamma_{67}/\Gamma_{44}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.0 \times 10^{-2}$	90	AUBERT	07U BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ K^+ \pi^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$  $\Gamma_{68}/\Gamma_{48}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.047 \pm 0.011 \pm 0.008</math></b>	105	ABE	02C BELL	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+ \pi^+ \pi^-)$  $\Gamma_{69}/\Gamma_{48}$ Unseen decay modes of the  $K^*(892)^0$  are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.078 \pm 0.018 \pm 0.013</math></b>	49	LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

 $\Gamma(\Sigma^- K^+ \pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$  $\Gamma_{70}/\Gamma_{69}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.35$	90	LINK	02G FOCS	$\gamma$ nucleus, $\approx 180$ GeV

**Doubly Cabibbo-suppressed modes** $\Gamma(p K^+ \pi^-)/\Gamma(p K^- \pi^+)$  $\Gamma_{71}/\Gamma_2$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.77 \pm 0.27</math> OUR AVERAGE</b>				Error includes scale factor of 1.9.
1.65 $\pm 0.15 \pm 0.05$	392	AAIJ	18V LHCb	$\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$
2.35 $\pm 0.27 \pm 0.21$	3379	YANG	16 BELL	At or near $\gamma$ s

**Semileptonic modes** $\Gamma(\Lambda e^+ \nu_e)/\Gamma_{\text{total}}$  $\Gamma_{72}/\Gamma$ 

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.63 \pm 0.38 \pm 0.20</math></b>	104	ABLIKIM	15Y BES3	$567 \text{ pb}^{-1}, 4.599 \text{ GeV}$

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(e^+ \text{anything})$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{72}/\Gamma_{74}$
<b>91.9±12.5±5.4</b>	214	ABLIKIM	18AF BES3	$e^+ e^-$ 4.6 GeV	

 $\Gamma(\Lambda e^+ \nu_e)/\Gamma(p K^- \pi^+)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{72}/\Gamma_2$
-------	-------------	------	---------	------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.43±0.08 1,2 BERGFELD 94 CLE2  $e^+ e^- \approx \gamma(4S)$

0.38±0.14 2,3 ALBRECHT 91G ARG  $e^+ e^- \approx 10.4$  GeV

<sup>1</sup> BERGFELD 94 measures  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.87 \pm 0.28 \pm 0.69)$  pb.

<sup>2</sup> To extract  $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)/\Gamma(\Lambda_c^+ \rightarrow p K^- \pi^+)$ , we use  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c \rightarrow p K^- \pi^+) = (11.2 \pm 1.3)$  pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (EVERY 91).

<sup>3</sup> ALBRECHT 91G measures  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.20 \pm 1.28 \pm 0.71)$  pb.

 $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{73}/\Gamma$
<b>3.49±0.46±0.27</b>	79	ABLIKIM	17D	BES3	$e^+ e^-$ at 4.6 GeV

 $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(p K^- \pi^+)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{73}/\Gamma_2$
-------	-------------	------	---------	------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.40±0.09 1,2 BERGFELD 94 CLE2  $e^+ e^- \approx \gamma(4S)$

0.35±0.20 2,3 ALBRECHT 91G ARG  $e^+ e^- \approx 10.4$  GeV

<sup>1</sup> BERGFELD 94 measures  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (4.43 \pm 0.51 \pm 0.64)$  pb.

<sup>2</sup> To extract  $\Gamma(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow p K^- \pi^+)$ , we use  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c \rightarrow p K^- \pi^+) = (11.2 \pm 1.3)$  pb, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (EVERY 91).

<sup>3</sup> ALBRECHT 91G measures  $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (3.91 \pm 2.02 \pm 0.90)$  pb.

 $\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda e^+ \nu_e)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{73}/\Gamma_{72}$
-------	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.96±0.16±0.04 <sup>1</sup> ABLIKIM 17D BES3  $e^+ e^-$  at 4.6 GeV

<sup>1</sup> This is the ratio of the ABLIKIM 17D  $\Lambda \mu^+ \nu_e$  branching fraction and the ABLIKIM 15Y  $\Lambda e^+ \nu_e$  branching fraction (see above), and so is not an independent measurement.

**Inclusive modes** $\Gamma(e^+ \text{anything})/\Gamma_{\text{total}}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{74}/\Gamma$
<b>3.95±0.34±0.09</b>	214	ABLIKIM	18AF BES3	$e^+ e^-$ 4.6 GeV	

$\Gamma(p \text{ anything})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.50±0.08±0.14</b>	<sup>1</sup> CRAWFORD 92	CLEO	$e^+ e^-$ 10.5 GeV

<sup>1</sup> This CRAWFORD 92 value includes protons from  $\Lambda$  decay. The value is model dependent, but account is taken of this in the systematic error.

 $\Gamma(n \text{ anything})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.50±0.08±0.14</b>	<sup>1</sup> CRAWFORD 92	CLEO	$e^+ e^-$ 10.5 GeV

<sup>1</sup> This CRAWFORD 92 value includes neutrons from  $\Lambda$  decay. The value is model dependent, but account is taken of this in the systematic error.

 $\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>38.2<sup>+2.8</sup><sub>-2.2</sub>±0.9</b>	700	ABLIKIM	18E BES3	$e^+ e^-$ at 4.6 GeV

 $\Gamma(3\text{prongs})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.24±0.07±0.04</b>	KAYIS-TOPAK.03	CHRS	$\nu_\mu$ emulsion, $\bar{E}=27$ GeV

**Rare or forbidden modes** $\Gamma(pe^+ e^-)/\Gamma_{\text{total}}$ 

A test for the  $\Delta C=1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.5 × 10<sup>-6</sup></b>	90	$4.0 \pm 7.1$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(p\mu^+ \mu^- \text{ non-resonant})/\Gamma_{\text{total}}$ 

A test for the  $\Delta C=1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;7.7 × 10<sup>-8</sup></b>	90	AAIJ	18N LHCb	Ratio to $p\phi$ , $\phi \rightarrow \mu^+ \mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4.4 \times 10^{-5}$	90	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$
$<3.4 \times 10^{-4}$	90	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV

 $\Gamma(pe^+ \mu^-)/\Gamma_{\text{total}}$ 

A test of lepton family-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;9.9 × 10<sup>-6</sup></b>	90	$-0.7 \pm 3.0$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(pe^- \mu^+)/\Gamma_{\text{total}}$ 

A test of lepton family-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;19 × 10<sup>-6</sup></b>	90	$6.2 \pm 4.9$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma(\bar{p}2e^+)/\Gamma_{\text{total}}$ 

A test of lepton- and baryon-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.7 × 10<sup>-6</sup></b>	90	$-1.5 \pm 4.5$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\bar{p}2\mu^+)/\Gamma_{\text{total}}$ 

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.4 \times 10^{-6}$	90	$0.0 \pm 2.2$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma_{84}/\Gamma$  $\Gamma(\bar{p}e^+\mu^+)/\Gamma_{\text{total}}$ 

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<16 \times 10^{-6}$	90	$10.1 \pm 6.8$	LEES	11G BABR	$e^+ e^- \approx \gamma(4S)$

 $\Gamma_{85}/\Gamma$  $\Gamma(\Sigma^-\mu^+\mu^+)/\Gamma_{\text{total}}$ 

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-4}$	90	0	KODAMA	95 E653	$\pi^-$ emulsion 600 GeV

 $\Gamma_{86}/\Gamma$  $\Lambda_c^+ \text{ DECAY PARAMETERS}$ 

See the note on "Baryon Decay Parameters" in the neutron Listings.

 $\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda\pi^+$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>-0.84 \pm 0.09</math> OUR AVERAGE</b>				
$-0.80 \pm 0.11 \pm 0.02$		ABLIKIM	19AX BES3	$e^+ e^-$ at 4.6 GeV
$-0.78 \pm 0.16 \pm 0.19$		LINK	06A FOCS	$\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV
$-0.94 \pm 0.21 \pm 0.12$	414	<sup>1</sup> BISHAI	95 CLE2	$e^+ e^- \approx \gamma(4S)$
$-0.96 \pm 0.42$		ALBRECHT	92 ARG	$e^+ e^- \approx 10.4$ GeV
$-1.1 \pm 0.4$	86	AVERY	90B CLEO	$e^+ e^- \approx 10.6$ GeV

<sup>1</sup> BISHAI 95 actually gives  $\alpha = -0.94^{+0.21+0.12}_{-0.06-0.06}$ , chopping the errors at the physical limit  $-1.0$ . However, for  $\alpha \approx -1.0$ , some experiments should get unphysical values ( $\alpha < -1.0$ ), and for averaging with other measurements such values (or errors that extend below  $-1.0$ ) should *not* be chopped.

 $\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^+\pi^0$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>-0.55 \pm 0.11</math> OUR AVERAGE</b>				
$-0.57 \pm 0.10 \pm 0.07$		ABLIKIM	19AX BES3	$e^+ e^-$ at 4.6 GeV
$-0.45 \pm 0.31 \pm 0.06$	89	BISHAI	95 CLE2	$e^+ e^- \approx \gamma(4S)$

 $\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^0\pi^+$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>-0.73 \pm 0.17 \pm 0.07</math></b>	ABLIKIM	19AX BES3	$e^+ e^-$ at 4.6 GeV

## $\alpha$ FOR $\Lambda_c^+ \rightarrow \Lambda \ell^+ \nu_\ell$

The experiments don't cover the complete (or same incomplete)  $M(\Lambda \ell^+)$  range, but we average them together anyway.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.86±0.04 OUR AVERAGE</b>				
-0.86±0.03±0.02	3201	<sup>1</sup> HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$
-0.91±0.42±0.25		<sup>2</sup> ALBRECHT	94B	ARG $e^+ e^- \approx 10$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.82 <sup>+0.09</sup> <sub>-0.06</sub> <sup>+0.06</sup> <sub>-0.03</sub>	700	<sup>3</sup> CRAWFORD	95	CLE2 See HINSON 05
-0.89 <sup>+0.17</sup> <sub>-0.11</sub> <sup>+0.09</sup> <sub>-0.05</sub>	350	<sup>4</sup> BERGFELD	94	CLE2 See CRAWFORD 95

<sup>1</sup>HINSON 05 measures the form-factor ratio  $R \equiv f_2/f_1$  for  $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$  events to be  $-0.31 \pm 0.05 \pm 0.04$  and the pole mass to be  $2.21 \pm 0.08 \pm 0.14$  GeV/c<sup>2</sup>, and from these calculates  $\alpha$ , averaged over  $q^2$ , where  $\langle q^2 \rangle = 0.67$  (GeV/c)<sup>2</sup>.

<sup>2</sup>ALBRECHT 94B uses  $\Lambda e^+$  and  $\Lambda \mu^+$  events in the mass range  $1.85 < M(\Lambda \ell^+) < 2.20$  GeV.

<sup>3</sup>CRAWFORD 95 measures the form-factor ratio  $R \equiv f_2/f_1$  for  $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$  events to be  $-0.25 \pm 0.14 \pm 0.08$  and from this calculates  $\alpha$ , averaged over  $q^2$ , to be the above.

<sup>4</sup>BERGFELD 94 uses  $\Lambda e^+$  events.

## $\alpha$ FOR $\Lambda_c^+ \rightarrow p K_S^0$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.18±0.43±0.14</b>	ABLIKIM	19AX BES3	$e^+ e^-$ at 4.6 GeV

## $\Lambda_c^+, \bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

### $(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} \pi^-$

This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.07±0.19±0.24</b>	LINK	06A FOCS	$\gamma$ A, $E_\gamma \approx 180$ GeV

### $(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$

This is zero if CP is conserved.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.00±0.03±0.02</b>	HINSON	05	CLEO $e^+ e^- \approx \gamma(4S)$

### $A_{CP}(\Lambda X)$ in $\Lambda_c \rightarrow \Lambda X, \bar{\Lambda}_c \rightarrow \bar{\Lambda} X$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1<sup>+7.0</sup><sub>-6.6</sub><sup>+1.6</sup></b>	700	ABLIKIM	18E BES3	$e^+ e^-$ at 4.6 GeV

### $\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow p K^+ K^-) - A_{CP}(\Lambda_c^+ \rightarrow p \pi^+ \pi^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>0.30±0.91±0.61</b>	<sup>1</sup> AAIJ	18R LHCb	$p p$ 7, 8 TeV

<sup>1</sup>AAIJ 18R applies phase-space-dependent weights to the  $\Lambda_c^+ \rightarrow p \pi^+ \pi^-$  sample to align its kinematics with the  $\Lambda_c^+ \rightarrow p K^+ K^-$  sample.

# $\Lambda_c^+$ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

AAIJ	19AG	PR D100 032001	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	19AX	PR D100 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19X	CP C43 083002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Y	PR D99 032010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18N	PR D97 091101	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18R	JHEP 1803 182	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18V	JHEP 1803 043	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	18AF	PRL 121 251801	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18E	PRL 121 062003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18Y	PL B783 200	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BERGER	18	PR D98 112006	M. Berger <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17D	PL B767 42	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17H	PRL 118 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17Q	PR D95 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17Y	PL B772 388	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PAL	17	PR D96 051102	B. Pal <i>et al.</i>	(BELLE Collab.)
ABLIKIM	16	PRL 116 052001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16U	PRL 117 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
YANG	16	PRL 117 011801	S.B. Yang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	15Y	PRL 115 221805	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZUPANC	14	PRL 113 042002	A. Zupanc <i>et al.</i>	(BELLE Collab.)
LEES	11G	PR D84 072006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
VAZQUEZ-JA... 08	PL B666 299	E. Vazquez-Jauregui <i>et al.</i>	(SELEX Collab.)	
AUBERT	07U	PR D75 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	06A	PL B634 165	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT,B	05S	PR D72 052006	B. Aubert <i>et al.</i>	(BABAR Collab.)
HINSON	05	PRL 94 191801	J.W. Hinson <i>et al.</i>	(CLEO Collab.)
LINK	05F	PL B624 22	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
CRONIN-HEN... 03	PR D67 012001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	
KAYIS-TOPAK..03	PL B555 156	A. Kayis-Topaksu <i>et al.</i>	(CERN CHORUS Collab.)	
ABE	02C	PL B524 33	K. Abe <i>et al.</i>	(KEK BELLE Collab.)
LINK	02C	PRL 88 161801	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02G	PL B540 25	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)
KUSHNR... 01	PL B524 5243	A. Kushnirenko <i>et al.</i>	(FNAL SELEX Collab.)	
MAHMOOD	01	PRL 86 2232	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
AITALA	00	PL B471 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ALAM	98	PR D57 4467	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96E	PRPL 276 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	96C	PR D53 1013	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95B	PL B342 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMMAR	95	PRL 74 3534	R. Ammar <i>et al.</i>	(CLEO Collab.)
BISHAI	95	PL B350 256	M. Bishai <i>et al.</i>	(CLEO Collab.)
CRAWFORD	95	PRL 75 624	G. Crawford <i>et al.</i>	(CLEO Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	94B	PL B326 320	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEEV	94	PAN 57 1370	A.N. Aleev <i>et al.</i>	(Serpukhov BIS-2 Collab.)
Translated from YF 57 1443.				
AVERY	94	PL B325 257	P. Avery <i>et al.</i>	(CLEO Collab.)
BERGFELD	94	PL B323 219	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	94E	PL B328 193	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	93	PRL 71 2391	P. Avery <i>et al.</i>	(CLEO Collab.)
BOZEK	93	PL B312 247	A. Bozek <i>et al.</i>	(CERN NA32 Collab.)
FRAEBETTI	93D	PRL 70 1755	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRAEBETTI	93H	PL B314 477	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KUBOTA	93	PRL 71 3255	Y. Kubota <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92	PL B274 239	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARLAG	92	PL B283 465	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
JEZABEK	92	PL B286 175	M. Jezabek, K. Rybicki, R. Rylko	(CRAC)
ALBRECHT	91G	PL B269 234	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	91	PR D43 3599	P. Avery <i>et al.</i>	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)

ALVAREZ	90B	PL B246 256	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90	PR D41 801	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AVERY	90B	PRL 65 2842	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	90D	ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRAEBETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BARLAG	89	PL B218 374	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
AGUILAR-...	88B	ZPHY C40 321	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B189 254	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		PL B199 462	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
Also		SJNP 48 833	M. Begalli <i>et al.</i>	(LEBC-EHS Collab.)
		Translated from YAF 48 1310.		
ALBRECHT	88C	PL B207 109	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88B	PRL 60 1379	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AMENDOLIA	87	ZPHY C36 513	S.R. Amendolia <i>et al.</i>	(CERN NA1 Collab.)
JONES	87	ZPHY C36 593	G.T. Jones <i>et al.</i>	(CERN WA21 Collab.)
BOSETTI	82	PL 109B 234	P.C. Bosetti <i>et al.</i>	(AACH3, BONN, CERN+)
BASILE	81B	NC 62A 14	M. Basile <i>et al.</i>	(CERN, BGNA, PGIA, FRAS)
CALICCHIO	80	PL 93B 521	M. Calicchio <i>et al.</i>	(BARI, BIRM, BRUX+)

## OTHER RELATED PAPERS

MIGLIOZZI	99	PL B462 217	P. Migliozzi <i>et al.</i>
DUNIETZ	98	PR D58 094010	I. Dunietz