

**$\psi(4160)$**

$I^G(J^{PC}) = 0^-(1^{--})$

### **$\psi(4160)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4191 ± 5 OUR AVERAGE</b>			
4191 + 9 - 8	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	<sup>1</sup> ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 ± 7	<sup>2</sup> MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4151 ± 4	<sup>3</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4155 ± 5	<sup>4</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02C) data.

### **$\psi(4160)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>70 ±10 OUR AVERAGE</b>			
65 +22 -16	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8±12.3	<sup>1</sup> ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 ±14	<sup>2</sup> MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±10	<sup>3</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±16	<sup>4</sup> SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
78 ±20	BRANDELIK	78C DASP	$e^+ e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02C) data.

## $\psi(4160)$ DECAY MODES

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle’s central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
$\Gamma_2 \mu^+ \mu^-$	seen	
$\Gamma_3 D\bar{D}$	seen	
$\Gamma_4 D^0 \bar{D}^0$	seen	
$\Gamma_5 D^+ D^-$	seen	
$\Gamma_6 D^* \bar{D} + \text{c.c.}$	seen	
$\Gamma_7 D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_8 D^*(2010)^+ D^- + \text{c.c.}$	seen	
$\Gamma_9 D^* \bar{D}^*$	seen	
$\Gamma_{10} D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
$\Gamma_{11} D^*(2010)^+ D^*(2010)^-$	seen	
$\Gamma_{12} D^0 D^- \pi^+ + \text{c.c. (excl.)}$ $D^*(2007)^0 \bar{D}^0 + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)$	seen	
$\Gamma_{14} D^0 D^{*-} \pi^+ + \text{c.c. (excl.)}$ $D^*(2010)^+ D^*(2010)^-$	not seen	
$\Gamma_{15} D_s^+ D_s^-$	not seen	
$\Gamma_{16} D_s^{*+} D_s^- + \text{c.c.}$	seen	
$\Gamma_{17} J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{18} J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%
$\Gamma_{19} J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%
$\Gamma_{20} J/\psi \eta$	$< 8 \times 10^{-3}$	90%
$\Gamma_{21} J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{22} J/\psi \eta'$	$< 5 \times 10^{-3}$	90%
$\Gamma_{23} J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{24} \psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{25} \chi_{c1} \gamma$	$< 5 \times 10^{-3}$	90%
$\Gamma_{26} \chi_{c2} \gamma$	$< 1.3 \%$	90%
$\Gamma_{27} \chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{28} \chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%
$\Gamma_{29} h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%
$\Gamma_{30} h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{31} h_c(1P) \eta$	$< 2 \times 10^{-3}$	90%
$\Gamma_{32} h_c(1P) \pi^0$	$< 4 \times 10^{-4}$	90%
$\Gamma_{33} \phi \pi^+ \pi^-$	$< 2 \times 10^{-3}$	90%

$\Gamma_{34}$	$\gamma\chi_{c1}(3872) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 6.8	$\times 10^{-5}$	90%
$\Gamma_{35}$	$\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.36	$\times 10^{-4}$	90%
$\Gamma_{36}$	$\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.18	$\times 10^{-4}$	90%
$\Gamma_{37}$	$\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-$	< 1.47	$\times 10^{-4}$	90%
$\Gamma_{38}$	$\gamma\chi_{c1}(3872) \rightarrow \gamma\gamma J/\psi$	< 1.05	$\times 10^{-4}$	90%
$\Gamma_{39}$	$\gamma X(3915) \rightarrow \gamma\gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%
$\Gamma_{40}$	$\gamma X(3930) \rightarrow \gamma\gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%
$\Gamma_{41}$	$\gamma X(3940) \rightarrow \gamma\gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%
$\Gamma_{42}$	$K^+K^-$			
$\Gamma_{43}$	$K_S^0 K^\pm \pi^\mp$			

### $\psi(4160)$ PARTIAL WIDTHS

#### $\Gamma(e^+e^-)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_1$
<b>0.48±0.22</b>	1 ABLIKIM	08D	BES2 $e^+e^- \rightarrow$ hadrons	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.4 to 1.1	2 MO	10	RVUE $e^+e^- \rightarrow$ hadrons	
0.83±0.08	3 SETH	05A	RVUE $e^+e^- \rightarrow$ hadrons	
0.84±0.13	4 SETH	05A	RVUE $e^+e^- \rightarrow$ hadrons	
0.77±0.23	BRANDELIK	78C	DASP $e^+e^-$	

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different  $e^+e^-$  partial widths. We quote only the range of values.

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>4</sup> From a fit to BES (BAI 02C) data.

#### $\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

#### $\Gamma(J/\psi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{22}\Gamma_1/\Gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.17±0.04	86	1,2 ABLIKIM	20A	BES3 $e^+e^- \rightarrow \eta' J/\psi$	
1.07±0.09	86	1,3 ABLIKIM	20A	BES3 $e^+e^- \rightarrow \eta' J/\psi$	

<sup>1</sup> Based on a fit to  $\sigma(e^+e^- \rightarrow \eta' J/\psi)$  from  $\sqrt{s} = 4.18$  to 4.60 GeV assuming interfering  $\psi(4160)$  and  $\psi(4260)$  contributions. At  $\sqrt{s} = 4.18$  GeV,  $\sigma(e^+e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$  pb.

<sup>2</sup> Solution I of the fit, corresponding to a phase of  $-0.03 \pm 0.44$  rad.

<sup>3</sup> Solution II of the fit, corresponding to a phase of  $2.54 \pm 0.04$  rad.

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{25}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.2	90	1 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{26}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.1	90	1 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{43}\Gamma_1/\Gamma$		
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.71 $\pm 0.13 \pm 0.12$	1 ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0095 $\pm 0.0088 \pm 0.0004$	2 ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Solution I of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

<sup>2</sup> Solution II of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

### $\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$		
<u>VALUE (units <math>10^{-8}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.8 $\pm 0.9 \pm 0.9$	1 WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$
12.8 $\pm 1.7 \pm 2.0$	2 WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

<sup>1</sup> Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

<sup>2</sup> Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

### $\psi(4160)$ BRANCHING RATIOS

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	1 AAIJ	13BC LHCb	$B^+ \rightarrow K^+\mu^+\mu^-$

<sup>1</sup> AAIJ 13BC report  $B(B^+ \rightarrow K^+\psi(4160)) B(\psi(4160) \rightarrow \mu^+\mu^-) = (3.5^{+0.9}_{-0.8}) \times 10^{-9}$ .

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	$\Gamma_3/\Gamma_9$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.02 $\pm 0.03 \pm 0.02$	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ VALUE**seen****seen**

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

DOCUMENT IDCRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^0\bar{D}^0$ PAKHLOVA 08 BELL  $e^+e^- \rightarrow D^0\bar{D}^0\gamma$  $\Gamma_4/\Gamma$ TECN $e^+e^- \rightarrow D^0\bar{D}^0$  $D^0\bar{D}^0\gamma$  $e^+e^- \rightarrow D^0\bar{D}^0\gamma$  $\Gamma(D^+D^-)/\Gamma_{\text{total}}$ VALUE**seen****seen**

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

DOCUMENT IDCRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^+D^-$ PAKHLOVA 08 BELL  $e^+e^- \rightarrow D^+D^-\gamma$  $\Gamma_5/\Gamma$ TECN $e^+e^- \rightarrow D^+D^-$  $D^+D^-\gamma$  $e^+e^- \rightarrow D^+D^-\gamma$  $\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$ VALUE**seen****seen**DOCUMENT IDAUBERT 09M BABR  $e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$ CRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^{*0}\bar{D}^0$  $\Gamma_7/\Gamma$ TECN $D^{*0}\bar{D}^0\gamma$  $D^{*0}\bar{D}^0$  $\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$ VALUE**seen****seen****seen**

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

DOCUMENT ID1 ZHUKOVA 18 BELL  $e^+e^- \rightarrow D^{*+}D^-\gamma$ AUBERT 09M BABR  $e^+e^- \rightarrow D^{*+}D^-\gamma$ CRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^{*+}D^-$  $\Gamma_8/\Gamma$ TECN $D^{*+}D^-\gamma$  $D^{*+}D^-$ <sup>1</sup> Supersedes PAHKLOVA 07. $\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$ VALUE**0.34±0.14±0.05**DOCUMENT IDAUBERT 09M BABR  $e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$  $\Gamma_6/\Gamma_9$ TECN $\gamma D^{(*)}\bar{D}^{(*)}$  $\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ VALUE**seen****seen**DOCUMENT IDAUBERT 09M BABR  $e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$ CRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$  $\Gamma_{10}/\Gamma$ TECN $D^{*0}\bar{D}^{*0}\gamma$  $D^{*0}\bar{D}^{*0}$  $\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$ VALUE**seen****seen****seen**

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

DOCUMENT ID1 ZHUKOVA 18 BELL  $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$ AUBERT 09M BABR  $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$ CRONIN-HEN..09 CLEO  $e^+e^- \rightarrow D^{*+}D^{*-}$  $\Gamma_{11}/\Gamma$ TECN $D^{*+}D^{*-}\gamma$  $D^{*+}D^{*-}$ <sup>1</sup> Supersedes PAHKLOVA 07.

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})) / \Gamma_{\text{total}}$	$\Gamma_{12}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>not seen</b>	PAKHLOVA    08A    BELL $e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)) / \Gamma_{\text{total}}$	$\Gamma_{13}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>seen</b>	CRONIN-HEN..09    CLEO $e^+ e^- \rightarrow D \bar{D}^* \pi$
$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-) / \Gamma_{\text{total}}$	$\Gamma_{14}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>not seen</b>	PAKHLOVA    09    BELL $e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$
$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}}$	$\Gamma_{15}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>not seen</b>	PAKHLOVA    11    BELL $e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	DEL-AMO-SA..10N    BABR $e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>not seen</b>	CRONIN-HEN..09    CLEO $e^+ e^- \rightarrow D_s^+ D_s^-$
$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}}$	$\Gamma_{16}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>seen</b>	PAKHLOVA    11    BELL $e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	DEL-AMO-SA..10N    BABR $e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
<b>seen</b>	CRONIN-HEN..09    CLEO $e^+ e^- \rightarrow D_s^{*+} D_s^-$
$\Gamma(J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}}$	$\Gamma_{17}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>&lt;3</b> 90	COAN    06    CLEO $4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$
$\Gamma(J/\psi \pi^0 \pi^0) / \Gamma_{\text{total}}$	$\Gamma_{18}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>&lt;3</b> 90	COAN    06    CLEO $4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$
$\Gamma(J/\psi K^+ K^-) / \Gamma_{\text{total}}$	$\Gamma_{19}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>&lt;2</b> 90	COAN    06    CLEO $4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$
$\Gamma(J/\psi \eta) / \Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b>&lt;8</b> 90	COAN    06    CLEO $4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>	
possibly seen	<sup>1</sup> ABLIKIM    15L    BES3 $e^+ e^- \rightarrow J/\psi \eta$
seen	WANG    13B    BELL $e^+ e^- \rightarrow J/\psi \eta \gamma$

<sup>1</sup> An enhancement around 4.2 GeV is observed.

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$				$\Gamma_{21}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(J/\psi\eta')/\Gamma_{\text{total}}$				$\Gamma_{22}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_{23}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$				$\Gamma_{24}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$				$\Gamma_{25}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<7	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$				$\Gamma_{26}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_{27}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_{28}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$				$\Gamma_{29}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

$\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$				$\Gamma_{30}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

$\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90		1 PEDLAR	11	CLEO $e^+ e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 41 2 ABLIKIM 17R BES3  $e^+ e^- \rightarrow h_c(1P)\eta$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+ e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

<sup>2</sup> An enhancement around 4.2 GeV is observed.

 $\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{32}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.4	90	1 PEDLAR	11	CLEO $e^+ e^- \rightarrow h_c(1P)\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+ e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{33}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(\gamma\chi_{c1}(3872) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{34}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
< $0.68 \times 10^{-4}$	90	1 XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
< $1.36 \times 10^{-4}$	90	1 XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{36}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
< $1.18 \times 10^{-4}$	90	1 XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{37}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
< $1.47 \times 10^{-4}$	90	1 XIAO	$\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma\chi_{c1}(3872) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
< $1.05 \times 10^{-4}$	90	1 XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{39}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$<1.26 \times 10^{-4}$	90	<sup>1</sup> XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{40}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$<0.88 \times 10^{-4}$	90	<sup>1</sup> XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{41}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
$<1.79 \times 10^{-4}$	90	<sup>1</sup> XIAO	$\psi(4160) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration. $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{42}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

 $\bullet \bullet \bullet$  We do not use the following data for averages, fits, limits, etc.  $\bullet \bullet \bullet$ 

$<2 \times 10^{-5}$	90	<sup>1</sup> DRUZHININ	15	RVUE $e^+ e^- \rightarrow \psi(3770)$
---------------------	----	------------------------	----	---------------------------------------

<sup>1</sup> DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes  $e^+ e^- \rightarrow K^+ K^-$  and  $e^+ e^- \rightarrow K_S^0 K_L^0$ . **$\psi(4160)$  REFERENCES**

ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan	
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PR L 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)