

**$K_0^*(1950)$** 

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the  $K^- \pi^+$  system. Needs confirmation. **$K_0^*(1950)$  MASS**

| <u>VALUE (MeV)</u>    | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u>                       |
|-----------------------|---|-------------|------------|--------------------------------------|
| <b>1945 ± 10 ± 20</b> | <sup>1</sup> ASTON  | 88          | LASS       | 0 11 $K^- p \rightarrow K^- \pi^+ n$ |
| • • •                 | We do not use the following data for averages, fits, limits, etc. • • • |             |            |                                      |
| 1917 ± 12             | <sup>2</sup> ZHOU   | 06          | RVUE       | $K p \rightarrow K^- \pi^+ n$        |
| 1820 ± 40             | <sup>3</sup> ANISOVICH  | 97C         | RVUE       | 11 $K^- p \rightarrow K^- \pi^+ n$   |

<sup>1</sup>We take the central value of the two solutions and the larger error given.<sup>2</sup>S-matrix pole. Using ASTON 88 and assuming  $K_0^*(700)$ ,  $K_0^*(1430)$ .<sup>3</sup>T-matrix pole. Reanalysis of ASTON 88 data. **$K_0^*(1950)$  WIDTH**

| <u>VALUE (MeV)</u>   | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u>                       |
|----------------------|---|-------------|------------|--------------------------------------|
| <b>201 ± 34 ± 79</b> | <sup>4</sup> ASTON  | 88          | LASS       | 0 11 $K^- p \rightarrow K^- \pi^+ n$ |
| • • •                | We do not use the following data for averages, fits, limits, etc. • • • |             |            |                                      |
| 145 ± 38             | <sup>5</sup> ZHOU   | 06          | RVUE       | $K p \rightarrow K^- \pi^+ n$        |
| 250 ± 100            | <sup>6</sup> ANISOVICH  | 97C         | RVUE       | 11 $K^- p \rightarrow K^- \pi^+ n$   |

<sup>4</sup>We take the central value of the two solutions and the larger error given.<sup>5</sup>S-matrix pole. Using ASTON 88 and assuming  $K_0^*(700)$ ,  $K_0^*(1430)$ .<sup>6</sup>T-matrix pole. Reanalysis of ASTON 88 data. **$K_0^*(1950)$  DECAY MODES**

| Mode                   | Fraction ( $\Gamma_i/\Gamma$ ) |
|------------------------|--------------------------------|
| $\Gamma_1$ $K^- \pi^+$ | (52 ± 14) %                    |

 **$K_0^*(1950)$  BRANCHING RATIOS**

| $\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$ | $\Gamma_1/\Gamma$   |  |
|---|---|--|
| <b>0.52 ± 0.08 ± 0.12</b>                 | <sup>7</sup> ASTON  |  |
| • • •                                     | We do not use the following data for averages, fits, limits, etc. • • •                 |  |
| ~ 0.60                                    | <sup>8</sup> ZHOU   |  |
|   | $K p \rightarrow K^- \pi^+ n$   |  |
|   | <sup>7</sup> We take the central value of the two solutions and the larger error given. |  |
|   | <sup>8</sup> S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$ , $K_0^*(1430)$ .  |  |

## $K_0^*(1950)$ REFERENCES

|           |     |             |   |
|-----------|-----|-------------|---|
| ZHOU      | 06  | NP A775 212 | Z.Y. Zhou, H.Q. Zheng                           |
| ANISOVICH | 97C | PL B413 137 | A.V. Anisovich, A.V. Sarantsev                  |
| ASTON     | 88  | NP B296 493 | D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS) |

---