

$\psi(4415)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

### $\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4421 ± 4 OUR ESTIMATE</b>			
<b>4415.1 ± 7.9</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4411 ± 7	<sup>3</sup> PAKHLOVA	08A BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4429 ± 9	<sup>5</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	$e^+e^-$
4414 ± 7	SIEGRIST	76 MRK1	$e^+e^-$

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>3</sup> Systematic uncertainties not estimated.

<sup>4</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>5</sup> From a fit to BES (BAI 02C) data.

### $\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>62 ± 20 OUR ESTIMATE</b>			
<b>71.5 ± 19.0</b>	<sup>6</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	<sup>7</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
77 ± 20	<sup>8</sup> PAKHLOVA	08A BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	<sup>9</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
118 ± 35	<sup>10</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C DASP	$e^+e^-$
33 ± 10	SIEGRIST	76 MRK1	$e^+e^-$

<sup>6</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .

<sup>7</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

<sup>8</sup> Systematic uncertainties not estimated.

<sup>9</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

<sup>10</sup> From a fit to BES (BAI 02C) data.

## $\psi(4415)$ DECAY MODES

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle’s central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $D\bar{D}$	seen	
$\Gamma_2$ $D^0\bar{D}^0$	seen	
$\Gamma_3$ $D^+D^-$	seen	
$\Gamma_4$ $D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_5$ $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_6$ $D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_7$ $D^*\bar{D}^*$	seen	
$\Gamma_8$ $D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen	
$\Gamma_9$ $D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen	
$\Gamma_{10}$ $D^0D^-\pi^+$ (excl. $D^*(2007)^0\bar{D}^0$ +c.c., $D^*(2010)^+D^-$ +c.c.)	$< 2.3$ %	90%
$\Gamma_{11}$ $D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+ + \text{c.c.}$	$(10 \pm 4)$ %	
$\Gamma_{12}$ $D^0D^{*-}\pi^+ + \text{c.c.}$	$< 11$ %	90%
$\Gamma_{13}$ $D_1(2420)\bar{D} + \text{c.c.}$	possibly seen	
$\Gamma_{14}$ $D_s^+D_s^-$	not seen	
$\Gamma_{15}$ $\omega\chi_{c2}$	possibly seen	
$\Gamma_{16}$ $D_s^{*+}D_s^- + \text{c.c.}$	seen	
$\Gamma_{17}$ $D_s^{*+}D_s^{*-}$	not seen	
$\Gamma_{18}$ $\psi_2(3823)\pi^+\pi^-$	possibly seen	
$\Gamma_{19}$ $\psi(3770)\pi^+\pi^-$	possibly seen	
$\Gamma_{20}$ $J/\psi\eta$	$< 6 \times 10^{-3}$	90%
$\Gamma_{21}$ $\chi_{c1}\gamma$	$< 8 \times 10^{-4}$	90%
$\Gamma_{22}$ $\chi_{c2}\gamma$	$< 4 \times 10^{-3}$	90%
$\Gamma_{23}$ $e^+e^-$	$(9.4 \pm 3.2) \times 10^{-6}$	

## $\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23}$
<b>0.58 ± 0.07 OUR ESTIMATE</b>					
<b>0.35 ± 0.12</b>		11 ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.4 to 0.8		12 MO	10 RVUE	$e^+e^- \rightarrow$ hadrons	
0.72 ± 0.11		13 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons	
0.64 ± 0.23		14 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons	
0.49 ± 0.13		BRANDELIK	78C DASP	$e^+e^-$	
0.44 ± 0.14		SIEGRIST	76 MRK1	$e^+e^-$	

- <sup>11</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .
- <sup>12</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different  $e^+e^-$  partial widths. We quote only the range of values.
- <sup>13</sup> From a fit to Crystal Ball (OSTERHELD 86) data.
- <sup>14</sup> From a fit to BES (BAI 02C) data.

### $\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{20}\Gamma_{23}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<3.6	90	WANG	13B	BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{21}\Gamma_{23}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.47	90	<sup>15</sup> HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

<sup>15</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{22}\Gamma_{23}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.3	90	<sup>16</sup> HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

<sup>16</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

### $\psi(4415)$ BRANCHING RATIOS

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
•••					We do not use the following data for averages, fits, limits, etc. •••
not seen		AUBERT	09M	BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		PAKHLOVA	08	BELL	$e^+e^- \rightarrow D^+D^-\gamma$
•••					We do not use the following data for averages, fits, limits, etc. •••
not seen		AUBERT	09M	BABR	$e^+e^- \rightarrow D^+D^-\gamma$

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$					$\Gamma_1/\Gamma_7$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.14±0.12±0.03</b>		AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	17 ZHUKOVA	18	BELL $e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	AUBERT	09M	BABR $e^+ e^- \rightarrow D^{*+} D^- \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	PAKHLOVA	07	BELL $e^+ e^- \rightarrow D^{*+} D^- \gamma$

<sup>17</sup> Supersedes PAKHLOVA 07. $\Gamma(D^* \bar{D} + \text{c.c.})/\Gamma(D^* \bar{D}^*)$   $\Gamma_4/\Gamma_7$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.17 ± 0.25 ± 0.03</b>	AUBERT	09M	BABR $e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

 $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AUBERT	09M	BABR $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

 $\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	18 ZHUKOVA	18	BELL $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	AUBERT	09M	BABR $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	PAKHLOVA	07	BELL $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

<sup>18</sup> Supersedes PAKHLOVA 07. $\Gamma(D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>10.5 ± 2.4 ± 3.8</b>	19 PAKHLOVA	08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

<sup>19</sup> Using  $4421 \pm 4$  MeV for the mass and  $62 \pm 20$  MeV for the width of  $\psi(4415)$ . $\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c.}, D^*(2010)^+ D^- + \text{c.c.})/\Gamma(D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$   $\Gamma_{10}/\Gamma_{11}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.22</b>	90	20 PAKHLOVA	08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

<sup>20</sup> Using  $4421 \pm 4$  MeV for the mass and  $62 \pm 20$  MeV for the width of  $\psi(4415)$ . $\Gamma(D^0 D^{*-} \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma \times \Gamma_{23}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.99 × 10<sup>-6</sup></b>	90	21 PAKHLOVA	09	BELL $e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

<sup>21</sup> Using  $4421 \pm 4$  MeV for the mass of  $\psi(4415)$ . $\Gamma(D_1(2420) \bar{D} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	22 ABLIKIM	19AR	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

<sup>22</sup> Evidence for  $e^+ e^- \rightarrow D_1(2420) \bar{D} + \text{c.c.}$  between  $\sqrt{s} = 4.3$  and 4.6 GeV, not necessarily resonant.

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$				$\Gamma_{14}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA...10N	BABR		$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

$\Gamma(\omega \chi_{c2})/\Gamma_{\text{total}}$				$\Gamma_{15}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
possibly seen	ABLIKIM	16A	BES3	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$				$\Gamma_{16}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA...10N	BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$				$\Gamma_{17}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
not seen	DEL-AMO-SA...10N	BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

$\Gamma(\psi(3770) \pi^+ \pi^-)/\Gamma_{\text{total}}$				$\Gamma_{19}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
possibly seen	<sup>23</sup> ABLIKIM	19AR	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

<sup>23</sup> Observe  $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$  at  $\sqrt{s} = 4.26, 4.36, \text{ and } 4.42$  GeV but cannot establish if continuum or resonant.

$\Gamma(\psi_2(3823) \pi^+ \pi^-)/\Gamma_{\text{total}}$				$\Gamma_{18}/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
possibly seen	19	<sup>24</sup> ABLIKIM	15S	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$

<sup>24</sup> From a fit of  $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$ ,  $\psi_2(3823) \rightarrow \chi_{c1} \gamma$  cross sections taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the  $\psi(4415)$  line shape.

## $\psi(4415)$ REFERENCES

ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...10N		PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)